



Research Article

FORMULATION AND EVALUATION OF ANTIAGING CREAM CONTAINING MANGIFERIN

Aimi Muneerah Shamsuddin, Mahendran Sekar *, Ahmad Zawawi Musa

Faculty of Pharmacy and Health Sciences, Universiti Kuala Lumpur, Royal College of Medicine Perak, Ipoh - 30450, Malaysia

*Corresponding Author Email: mahendransekar@unikl.edu.my

Article Received on: 26/06/18 Approved for publication: 30/06/18

DOI: 10.7897/2230-8407.09689

ABSTRACT

Skin aging can be described as irregular pigmentation, increased wrinkling, loss of elasticity, dryness and roughness. *Phaleria macrocarpa* is known as *Mahkota Dewa* in Malay and commonly available in Malaysia. The fruits contain large quantity of mangiferin which possess medicinal benefits including anti-diabetic, antioxidant, anti-proliferative, immunomodulatory, cardioprotective and diuretic properties. However, there is no formulation has been developed though it was reported for many biological properties. The aim of our study is to formulate and evaluate anti-aging cream containing mangiferin as an active ingredient. Two formulations with different concentrations of mangiferin were formulated by mixing oil and aqueous phase using homogenizer. Both the formulations showed significant antioxidant activity and tyrosinase inhibition. The formulated creams were consistent in quality and safe to be used on the skin. These results demonstrate that mangiferin isolated from *P. macrocarpa* has a good potential for cosmetic product development.

Keywords: *Phaleria macrocarpa*; Mangiferin; Antiaging; Antioxidant; Tyrosinase inhibition

INTRODUCTION

Skin aging is a complex process induced by constant exposure to ultraviolet (UV) irradiation and damages human skin. Melanogenesis is the complex process which is responsible for the formation of pigment melanins by melanocytes while skin wrinkling is associated with collagen deficiency resulting from reactive oxygen species that is generated from UV¹. Skin aging is associated with many factors and when doing researches on the topic, one will definitely come across to terms like ROS (reactive oxygen species) which generally a free radical that contributes to skin aging. ROS is basically generated during cellular metabolism, but the excessive production of ROS can lead to OS (oxidative stress). OS can be a precursor to a lot of health damaging issues including skin aging².

Phaleria macrocarpa is traditionally known as *Mahkota Dewa* that grows in tropical countries including Malaysia. This plant is called *simalakama* in Sumatra (Malay) and *Depok* (West Java, Indonesia). *P. macrocarpa* is a well-known medicinal plant and extensively used in traditional medicine. The fruits are used for treating various diseases like hypertension, kidney disorder, cancer and diabetes^{3,4}. Studies have also found that some beneficial bioactivities of this plant include anti-cancer, antioxidant, anti-cancer anti-bacterial, anti-hyperglycemic and anti-hyperlipidemia⁵. Mangiferin is a pharmacologically active compound present large amount in *P. macrocarpa*. Mangiferin showed antidiabetic⁶, anticancer⁷⁻⁸, antioxidant⁹, anti-inflammatory¹⁰, antiviral¹¹ and antimicrobial¹² activities.

However, so far there is no formulation have been developed though it was reported for potent antioxidant properties. Hence in the present study we are interested to formulate an antiaging cream using mangiferin obtained from *P. macrocarpa*. The tyrosinase inhibition activity and antioxidant activities of the

formulated creams were tested to check the potency of anti-aging effect.

MATERIALS AND METHODS

Collection and Extraction of *P. macrocarpa* Fruits

P. macrocarpa fruits were collected from Kg Paya Resak, Terengganu, Malaysia and authenticated by a botanist. The fruits were cut into small slices and shade dried. Then, they were grinded into coarse powder using blender. 250 g of plant powder was extracted with 1.5 L of methanol using soxhlet apparatus for 18-20 hours. The extract obtained then was concentrated in vacuo by rotary evaporation at reduced pressure and controlled temperature. After evaporation, the concentrated extract was kept in refrigerator and stored at 4°C until further use (Yield 5.5 g, 2.2%).

Isolation of Mangiferin

The concentrated mass (5.5 g) was suspended in 50 ml of 50% of ethanol then partitioned with 100 ml of dichloromethane for 4 times. The aqueous ethanolic phase was then hydrolysed by reflux with 2N sulphuric acid at pH 3 for an hour with continuous stirring. After cooled to room temperature, it was partitioned with ethyl acetate for 3 times. Subsequently, the ethyl acetate layer was combined and dried at 40 °C using vacuum rotary evaporator. The dried ethyl acetate fractions were dissolved in ethanol and left in a refrigerator overnight. After that, the precipitate was isolated by filtration. The crystallization was done by dissolving in aqueous ethanolic solution and again left in refrigerator overnight. Lastly, pale yellow needle-shaped crystals of mangiferin was isolated (2.15 g, 0.86%) and dried¹³. It was found to be homogenous by HPTLC when separated using the solvent system ethyl acetate: methanol: water: formic acid (6:2:1:1, R_f = 0.76). It was

characterized by comparing its melting point, IR, NMR and MS data with literature values¹⁴. The structure of mangiferin was shown in Fig. 1.




Fig. 1. Structure of Mangiferin

Formulation of Antiaging Cream Containing Mangiferin

The oily phase and aqueous phase were heated up to 70°C and mixed using homogenizer by addition of methyl of methyl paraben, mangiferin and fragrance. With constant mixing, remaining distilled water was added and stirring was continuing until mixture cools [15]. Cream was formed when the consistency of the mixture was viscous and opaque. The composition of formulations 1 and 2 were shown in Table 1.

Table 1. Composition of Antiaging Cream

Ingredients	Composition of formulations (%w/w)	
	F1	F2
Active Ingredient		
Mangiferin	1%	3%
Oily Phase		
Stearic acid	10%	10%
Cetyl alcohol	6%	6%
Liquid paraffin	6.6%	6.6%
Aqueous Phase		
Glycerin	5%	5%
Methyl paraben	0.05%	0.05%
Propylene glycol	30%	30%
Distilled water	q.s.	q.s.

The study is carried out as per International conference of Harmonization-Good Clinical Practices Guidelines (ICH-GCP) or as per Declaration of Helsinki guidelines.

Evaluation of Antiaging Cream¹⁵⁻¹⁷

The following parameters were used to evaluate the antiaging cream. The standard procedure was followed to evaluate all the parameters.

Determination of Type of Emulsion (Dye Method)

A scarlet red dye was mixed with the cream. A drop of the cream was placed on a microscopic slide and examined under a microscope. If the disperse globules appear red the continuous phase colourless, the cream is water-in-oil (w/o) type. The reverse condition is occurring in oil-in-water (o/w) type cream i.e. the disperse globules appear colourless and the continuous phase red.

pH of the Cream: The pH meter was calibrated using standard buffer solution. About 0.5 g of the cream was weighed and dissolved in 50 ml of distilled water and its pH was measured.

Homogeneity: The formulation was tested for homogeneity by visual appearance and touch.

Appearance: The appearance of the cream was judged by its color, pearlescence and roughness and graded.

After Feel: Emolliency, slipperiness and amount of residue left after the application of fixed amount of cream were checked.

Type of Smear: After application of cream, the type of film or smear formed on the skin was checked.

Removal: The ease of removal of the cream applied were examined by washing the applied part with tap water.

Irritancy test: An area (1sq.cm) was marked on the left-hand dorsal surface of human volunteers. The cream was applied to the specified area and time is noted. Presence of irritancy, erythema and edema were checked at regular intervals up to 24 h and reported.

Accelerated Stability Testing: Creams were divided into four parts and stability test was performed at 8°C ± 0.1°C in refrigerator and at 25°C ± 1°C, 40°C ± 1°C and 40°C ± 1°C in incubator with 75% relative humidity (RH), and the above parameters were observed for 8 weeks at weekly intervals.

In vitro antioxidant activity by DPPH radical scavenging method

A 100 µl aliquot of the different concentrations of mangiferin and its formulations along with standards were added to 2 ml of DPPH in methanol solution (100 µM) and incubated at 37°C for 20 min. After that, the absorbance of each solution was determined at 490 nm using UV-visible spectrophotometer¹⁸. The percentage inhibition was calculated as follows

$$\text{Percentage inhibition} = \frac{[(\text{Abs Control} - \text{Abs Sample}) \times 100]}{(\text{Abs control})}$$

Tyrosinase Inhibition Activity

Different concentrations of mangiferin and its formulations were assayed for tyrosinase inhibition by measuring their effect on tyrosinase activity. The reaction was carried out in 100 mM sodium phosphate buffer (pH 6.7) containing 1 mM L-tyrosine and 80 unit/mL mushroom tyrosinase at 37°C. The reaction mixture was pre-incubated for 10 min before adding mangiferin. The change of the absorbance at 475 nm (sometimes 490 nm) was measured¹⁵. The percent inhibition of tyrosinase was calculated as follows:

$$\text{Percentage inhibition} = \frac{[(\text{Abs Control} - \text{Abs Sample}) \times 100]}{(\text{Abs control})}$$

RESULTS AND DISCUSSION

Mangiferin was isolated from the fruits of *P. macrocarpa*, found to be homogenous by HPTLC (Fig. 2) when separated using the solvent system ethyl acetate: methanol: water: formic acid (6:2:1:1, $R_f = 0.76$). Obtained as pale-yellow needle shaped crystals, mp 270-272°C; yield 2.15 g, 0.86%; IR ν_{max} (KBr) cm^{-1} : 3365 (OH), 1651 (C=O), 1621 and 1594 (aromatic C=C), 1200 and 1096 (C-O), 1050, 828, 585 (Fig. 3); ^1H NMR (400 MHz, DMSO) δ : see Table 2; negative ESI-MS: m/z calculated for 422.08, Found: 444 for $[\text{M} + \text{Na} - \text{H}]$.




Fig. 2. HPTLC Spectrum of Mangiferin

Table 2. ¹H NMR Spectrum of Mangiferin

Proton	Literature Value ¹⁴	MGN
1-OH	13.76 (1H)	13.50 (1H)
6-OH	10.55 (1H)	10.80 (1H)
7-OH	10.55 (1H)	10.80 (1H)
3-OH	9.86 (1H)	9.80 (1H)
3',4'-OH	4.86 (2H)	4.90 (2H)
6'-OH	4.49 (1H)	4.49 (1H)
2'-OH	3.87 (1H)	4.60 (1H)
8-H	7.38 (1H)	7.40 (1H)
5-H	6.86 (1H)	6.90 (1H)
4-H	6.37 (1H)	6.40 (1H)
1'-H	4.60 (1H)	4.60 (1H)
2'-H	4.05 (1H)	4.05 (1H)
6'a-H	3.69 (1H)	3.69 (1H)
6'b-H	3.42 (1H)	3.41 (1H)
3'-H	3.18 (1H)	3.18 (1H)
4'-H	3.18 (1H)	3.18 (1H)
5'-H	3.18 (1H)	3.18 (1H)
Total number of protons	18	18

Table 5. *In-vitro* Antioxidant Activity by DPPH Method

Concentration (µg/ml)	% inhibition		
	Mangiferin	Formulation 1	Formulation 2
1000		51.42 ± 3.73	68.67 ± 1.57
500		32.35 ± 2.58	45.39 ± 3.52
250		23.52 ± 1.80	37.76 ± 1.74
125		--	25.45 ± 2.00
62.5	92.28 ± 2.92	--	19.97 ± 2.06
31.25	91.39 ± 1.13	--	16.35 ± 1.85
15.60	89.25 ± 1.00	--	--
7.80	81.45 ± 5.91	--	--
3.90	65.05 ± 9.98	--	--
1.95	47.65 ± 3.06	--	--
Ascorbic Acid (10 µg/ml)	92.45 ± 3.82		
Rutin (10 µg/ml)	48.25 ± 2.50		

-- No inhibition. Values are expressed as mean±SD (n=3).

Evaluation of Formulated Antiaging Cream

The dye test was confirming that all the formulations were o/w type of emulsion cream. The pH of the formulated cream was found to be in range of 4.80 to 5.60 which is good and recommended pH for the skin. Irritancy test was conducted with 5 healthy volunteers to identify the safety, skin irritation and allergic sensitization were scarce or absent. Both the formulations don't show redness, edema, inflammation and irritation during irritancy studies. The results indicate that both the formulations were safe to be used on the skin (Table 3).




Fig. 3. IR Spectrum of Mangiferin

Table 3: Type of Adverse Effect of Formulations

Formulation	Irritant	Erythema	Edema
F1	Nil	Nil	Nil
F2	Nil	Nil	Nil

Table 4. Physicochemical Evaluation of Formulated Cream

Parameter	Formulation 1	Formulation 2
Homogeneity	Good	Good
Appearance	No change in colour	No change in colour
Odour	Good	Good
Spreadability	Good	Good
After feel	Emollient	Emollient
Type of smear	Non greasy	Non greasy
Removal	Easy	Easy
Stability	Stable for 3 months	Stable for 3 months

Table 6. Tyrosinase Inhibition of Mangiferin Formulations

Concentration (µg/ml)	% inhibition		
	Mangiferin	Formulation 1	Formulation 2
1000	93.40 ± 2.10	54.49 ± 7.70	75.65 ± 2.53
500	71.65 ± 1.48	45.63 ± 2.68	64.67 ± 0.85
250	64.65 ± 4.04	37.35 ± 0.90	61.25 ± 1.98
125	56.93 ± 3.50	33.00 ± 2.85	57.57 ± 2.38
62.5	44.45 ± 3.67	20.73 ± 2.66	44.53 ± 2.55
31.25	32.13 ± 1.78	--	24.63 ± 3.90
15.60	25.42 ± 2.69	--	12.30 ± 1.15
7.80	14.67 ± 2.55	--	--
3.90	--	--	--
1.95	--	--	--
Gallic Acid (25 µg/ml)	52.35 ± 3.25		

-- No inhibition. Values are expressed as mean±SD (n=3).

The formulated antiaging creams were evaluated for several physicochemical tests and the results were shown in Table 4. The type of smear formed on the skin was not greasy after the application of all formulated creams. Both the formulated creams (F1 & F2) when applied on skin were easily removable by washing with water. All the formulations were producing a uniform distribution of extracts in the cream. This was confirmed by visual examination and by touch. When formulation kept for a long time, it was found that there were no changes in the colour of the cream. After feel test showed that the formulated creams were emollient and slipperiness.

All the physiochemical parameters were maintained during the accelerated stability studies at temperatures $8^{\circ}\text{C} \pm 0.1^{\circ}\text{C}$ in refrigerator and at $25^{\circ}\text{C} \pm 1^{\circ}\text{C}$, $40^{\circ}\text{C} \pm 1^{\circ}\text{C}$ and $40^{\circ}\text{C} \pm 1^{\circ}\text{C}$ in incubator for 12 weeks. The results of accelerated stability test showed that there were not any particular changes in the colour of the cream.

In-vitro Antioxidant Activity

Mangiferin showed potent antioxidant activity with percentage of inhibition value 92.28 ± 2.92 at $62.5 \mu\text{g/ml}$ in DPPH method. The formulation 1 and 2 showed potent antioxidant activity with percentage of inhibition value 51.42 ± 3.73 and 68.67 ± 1.57 at $1000 \mu\text{g/ml}$, respectively. Formulation 2 showed higher percentage of inhibition when compared to Formulation 1 in all the tested concentrations. However, standard ascorbic acid and rutin showed higher inhibition at low concentration when compared to mangiferin and its formulations. The results were showed in Table 5.

Tyrosinase Inhibition

Mangiferin showed potent tyrosinase inhibition activity with percentage of inhibition value 93.40 ± 2.10 at $1000 \mu\text{g/ml}$. The formulation 1 and 2 showed potent tyrosinase inhibition with percentage of inhibition value 54.49 ± 7.70 and 75.65 ± 2.53 at $1000 \mu\text{g/ml}$, respectively. Formulation 2 showed higher percentage of inhibition when compared to Formulation 1 in all the tested concentrations. However, standard gallic acid showed higher inhibition at low concentration when compared to mangiferin and its formulations. The results were showed in Table 6.

The use of antioxidants for a particular topical formulation appears to be an interesting approach to protect skin against oxidative stress caused by different extrinsic agents. To ensure the effectiveness of antioxidants against free radicals, it is essential to stabilize the final formulation as antioxidants are very unstable and can easily oxidize, becoming inactive before reaching its site of action¹⁵.

Our earlier laboratory studies reported that mangiferin showed potent antioxidant activity in DPPH, ABTS, H_2O_2 and nitric oxide methods¹⁸. It is beneficial effects on the process of skin aging, skin sun protection or skin cancer. There are many studies were reported that an acute exposure of human skin to UV radiation *in vivo* leads to oxidation of cellular biomolecules and that could be prevented by a prior antioxidant treatment. Hence, there is an increased demand for herbal cosmetics in the Malaysian markets. Therefore, the present study was tried to formulate an antiaging cream using mangiferin as an active ingredient.

The formulated cream was o/w type emulsion, hence can easily washed with water and gives better consumer compliance. Our study indicated that, both the formulated creams were stable with no signs of breakdown of emulsion and change in colour of the product. Also maintained constant pH, homogeneity emollient properties; they were not greasy and easily removable after the application. The irritancy test showed that there no severe erythema was occurred in any of the volunteers for formulation containing mangiferin. This indicates that the formulated creams were safe to the consumers.

CONCLUSION

Both the mangiferin formulations showed potent antioxidant activities and tyrosinase inhibition. The results demonstrated that

the formulated antiaging creams are safe and usable for the skin and having good potential for cosmetic product development.

REFERENCES

1. Rasul A, Akhtar N, Khan BA, Mahmood T, Uz Zaman S, Khan HM. Formulation development of a cream containing fennel extract: in vivo evaluation for anti-aging effects. *Pharmazie* 2012;67:54-58.
2. Khimara N, Mark ABM. Oxidative stress and ageing: The influence of environmental pollution, sunlight and diet on skin. *Cosmetics* 2017;4:1-8.
3. Tjandrawinata RR, Arifin PF, Tandrasasmita OM, Rahmi D and Aripin A. DLBS1425, a *Phaleria macrocarpa* (Scheff.) Boerl. extract confers antiproliferative and proapoptosis effects via eicosanoid pathway. *J Exp Ther Oncol*. 2010;8:187-201.
4. Winarno H and Katrin WE. Benzophenone glucoside isolated from the ethyl acetate of the bark of mahkota dewa (*Phaleria macrocarpa* (Scheff.) Boerl.) and its inhibitory activity on leukemia L1210 cell line. *Indo J Chem*. 2009;9:142-145.
5. Ramdani ED, Marlupi UD, Sinambela J, Tjandrawinata RR. Isolation and identification of compounds from *Phaleria macrocarpa* (Scheff.) Boerl fruit extract. *Asian Pacific J Trop Biomed* 2017;7:300-305.
6. Muruganandan S, Gupta S, Kataria M, Lal J, Gupta PK. Mangiferin protects the streptozotocin induced oxidative damage to cardiac and renal tissues in rats. *Toxicology*. 2002;176:165-73.
7. Rajendran P, Ekambaram G, Sakthisekaran D. Protective role of mangiferin against Benzo (a) pyrene induced lung carcinogenesis in experimental animals. *Biol Pharm Bull*. 2008;31:1053-8.
8. Leiro JM, Alvarez E, Arranz JA, Siso IG, Orallo F. *In vitro* effects of mangiferin on superoxide concentrations and expression of the inducible nitric oxide synthase, tumour necrosis factor- α and transforming growth factor- β genes. *Biochem Pharmacol*. 2003;65:1361-71.
9. Andreu GP, Delgado R, Velho JA, Curti C, Vercesi AE. Iron complexing activity of mangiferin, a naturally occurring glucosylxanthone, inhibits mitochondrial lipid peroxidation induced by Fe^{2+} -citrate. *Eur J Pharmacol*. 2005;513:47-55.
10. Garcia D, Delgado R, Ubeira FM, Leiro J. Modulation of rat macrophage function by the *Mangifera indica* L. extracts Vimang and mangiferin. *Int Immunopharmacol* 2002;2:797-806.
11. Zhu XM, Song JX, Huang ZZ, Wu YM, Yu MJ. Antiviral activity of mangiferin against herpes simplex virus type 2 *In vitro*. *Zhongguo Yao Li XueBao* 1993;14:452-4.
12. Bairy I, Reja S, Siddharth, Rao PS, Bhat M, Shivananda PG. Evaluation of antibacterial activity of *Mangifera indica* on anaerobic dental microflora based on *in vivo* studies. *Indian J Pathol Microbiol*. 2002;45:307-10.
13. Bhuvaneswari K. Isolation of mangiferin from leaves of *Mangifera indica* L. var alphonso. *Asian J Pharm Clin Res* 2013;6:173-174.
14. Mahendran, S. Molecules of Interest – Mangiferin - A Review. *Annual Res Rev Biol* 2014;5:307-320.
15. Mahendran S, Pavitra S, Afzan M. Formulation and Evaluation of Novel Antiaging Cream Containing Rambutan Fruit Extract. *Int J Pharma Sci Res* 2017;8:1056-1065.
16. Aswal A, Kalra H, Rout A. Preparation and evaluation of polyherbal cosmetic cream. *Scholars Res Library* 2013; 5:83-88.
17. Sabale V, Kunjwani H, Sabale P. Formulation and in vitro evaluation of the topical antiageing preparation of the fruit of

Benincasa hispida. J Ayurveda and Integ Med 2011; 2:124-128.

18. Mahendran S, Badami S, Ravi S, Thippeswamy BS, Veerapur VP. Synthesis and Evaluation of Analgesic and Anti-Inflammatory Activities of Most Active Free Radical Scavenging Derivatives of Mangiferin. British J Applied Sci Tech 2014;4:4959-4973.

Cite this article as:

Aimi Muneerah Shamsuddin et al. Formulation and evaluation of antiaging cream containing Mangiferin. Int. Res. J. Pharm. 2018;9(6):55-59 <http://dx.doi.org/10.7897/2230-8407.09689>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.