



## Research Article

### SYNTHESIS, CHARACTERIZATION, AND ANTIBACTERIAL ACTIVITY OF SCHIFF BASES DERIVED FROM THIOSEMICARBAZIDE, 2-ACETYL THIOPHENE AND THIOPHENE-2 ALDEHYDE

Chandra Mohan<sup>1\*</sup>, Vinod Kumar<sup>1</sup>, Sarla Kumari<sup>2</sup>

<sup>1</sup>Asst. Professor, K. R. Mangalam University, Gurgaon 122103, Haryana, India

<sup>2</sup>Department of Chemistry, S. D. Government College, Beawar 305901, Rajasthan, India

\*Corresponding Author Email: chandra.mohan@krmangalam.edu.in

Article Received on: 19/05/18 Approved for publication: 02/07/18

DOI: 10.7897/2230-8407.097141

#### ABSTRACT

Two Schiff bases 2-acetylthiophene thiosemicarbazone ( $L_1$ ) and thiophene-2-aldehyde thiosemicarbazone ( $L_2$ ) were prepared in excellent yield via the condensation of 2-acetyl thiophene/ thiophene-2-aldehyde and thiosemicarbazide. The lead (II) and nickel (II) metal complexes of the Schiff base ligands ( $[Pb(L_1)_2]Cl_2$  and  $[Ni(L_1)_2]Cl_2$ ) were also prepared. The ligands and complexes were tested against gram- negative [*E. Coli* (MTCC No: 452)] and gram- positive bacterial strains *Bacillus sp* [(MTCC No: 297)] by using Kirby-Bauer's method. The compound ( $[Pb(L_1)_2]Cl_2$ ) was found out to be more active against both bacterial strain (*E. coli* & *Bacillus sp*) having zone of inhibition  $12.45 \pm 1.2$  and  $11.27 \pm 1.1$  mm respectively.

**Keywords:** Antibacterial activity, Azomethine linkage, Schiff bases, Thiosemicarbazide

#### INTRODUCTION

Compound containing an azomethane group (-CH=N-) are known as Schiff bases. They are condensation products of ketones (or) aldehydes (aldehyde and ketones) with primary amines and were first reported by Hugo Schiff<sup>1</sup> in 1864. Schiff bases have wide applications in food industry, dye industry, analytical chemistry, catalysis, fungicidal, agrochemical and biological activities<sup>2</sup>. In recent years, the attention of Schiff bases and their metal complexes are increasing due to their remarkable biological and pharmacological applications<sup>3</sup>.

Metal complexes of Schiff bases as ligand with transition metals have received attention because of their biological and electrochemical activity including antitumor, antibacterial, fungicidal and anticarcinogenic properties<sup>4</sup>. The high affinity of the Schiff bases towards chelation with metal ions makes it suitable to for solid metal complexes<sup>5</sup>. They serve as models for biologically important species.

Many biologically important Schiff bases have been reported which possess antibacterial<sup>6-8</sup>, antifungal<sup>9-10</sup>, antimicrobial<sup>11-13</sup> and anti-HIV<sup>14, 15</sup> activities. The Schiff bases show the highest degree of hydrolysis and solubility in water at pH 5. Antitumor activity of the Schiff bases towards acidic tumors also increases considerably with the slight increase in watersolubility<sup>16</sup>. Several studies have also revealed that by the condensation of aldehyde or ketone with different heterocyclic compound, Schiff bases with potent antibacterial and antifungal activity was obtained<sup>17</sup>.

In the present work we have synthesized new Schiff bases, 2-acetylthiophene thiosemicarbazone ( $L_1$ ) and thiophene-2-aldehyde thiosemicarbazone ( $L_2$ ) and their Co(II), Cu(II), Zn(II) and Ni(II) complexes. The Schiff bases and their complexes were tested for antibacterial activity.

#### MATERIALS AND METHODS

All the chemicals used were of analytical grade (AR) and of the highest purity available. 2-acetyl thiophene, thiophene-2-aldehyde and thiosemicarbazide were procured from CDH and SRL. Metal salts were purchased from E. Merck and were used as received.

The C, H, N and S were analyzed on a Vario Micro Cube elemental analyzer, Model Vario-III. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker, Model DPX-300 NMR spectrophotometer using DMSO as solvent. The IR spectra were recorded as KBr pellets on a Perkin Elmer FT-IR spectrophotometer, Model No. BX-2.

#### Preparation of Schiff base Complexes<sup>18,19</sup>

##### Synthesis of ligand ( $L_1$ )

Ethanol solution of 2-acetyl thiophene (1.26g, 0.01 mol) was added in hot ethanolic solution (20 mL) of thiosemicarbazide (0.91g, 0.01 mol) with few drops of acetic acid with constant stirring. The mixture was refluxed at ~80 °C for 2 hours. On cooling, a light yellow coloured compound was separated out. It was then filtered, washed several times with 50% ethanol, dried and recrystallized from methanol (Fig. 1). Yield (88.0%), Melting Point: 140 °C.

##### Synthesis of ligand ( $L_2$ )

Thiophene-2-aldehyde thiosemicarbazone was synthesized by adding ethanolic solution of thiophene-2-aldehyde (1.12g, 0.01 mol) in hot ethanolic solution (20 mL) of thiosemicarbazide (0.91g, 0.01 mol) with few drops of acetic acid with constant stirring. The mixture was refluxed at ~80 °C for 3 hrs. On cooling, off-white coloured compound was separated out. It was filtered,

washed several times with 50% ethanol, dried and recrystallized from methanol (Fig. 2). Yield (75.0%), Melting Point: 160 °C.

#### Synthesis of metal complex

A general method was used for the synthesis of the complexes. The metal complexes of the Schiff bases, L<sub>1</sub> and L<sub>2</sub>, were prepared by adding a hot aqueous solution (20 mL) of corresponding metal ion (0.01M) dropwise with constant stirring to the hot ethanolic solution (0.02 M) of 2-acetylthiophene thiosemicarbazone/ thiophene-2-aldehyde thiosemicarbazone. The mixture was refluxed for 4-5 hours at 70 to 75 °C for complete complexation. On cooling a metal complex precipitate was separated out. The same was filtered, washed with 50% ethanol, dried over anhydrous calcium chloride and recrystallized from methanol (Fig. 3).

#### Biological Activity

The synthesized Schiff bases were screened for antibacterial activity.

#### Antibacterial Screening

All compounds including Schiff bases (L<sub>1</sub> & L<sub>2</sub>) and synthesized metal complexes of ligands L<sub>1</sub> i.e. ([Pb(L<sub>1</sub>)<sub>2</sub>]Cl<sub>2</sub> and [Ni(L<sub>1</sub>)<sub>2</sub>]Cl<sub>2</sub>) were analyzed against gram negative [*E. Coli* (MTCC No: 452)] and gram positive bacterial strains *Bacillus sp* [(MTCC No: 297)] through Kirby-Bauer's method<sup>20</sup>. Bacterial strains were cultivated and maintained by using Luria Bertani agar media (Hi-media laboratory, India). All compounds were tested at the concentration of 50 µg/ml against positive control (Ampicillin, 10 µg/ml) (Fig. 4 & 5).

#### Statistical Analysis

Values of antimicrobial assay were expressed as mean ± standard deviation (SD) and analyzed by using one-way analysis of variance (ANOVA) followed by Tukey's test. The value of  $p < 0.05$  was considered as statistically significant.

## RESULTS AND DISCUSSION

The Schiff base ligands and their metal complexes were characterized by modern spectroscopic methods such as FT-IR, UV-Vis., NMR and C, H, N, S elemental studies. Pre-confirmatory test for the synthesis of ligand was done through UV-Vis. and IR spectra. The composition of compound was confirmed by <sup>1</sup>H NMR, and CHN analysis.

#### Characterization of Schiff bases<sup>21</sup>

**2-acetylthiophene thiosemicarbazone (L<sub>1</sub>)-** Analytical calculation (%) for = C<sub>7</sub>H<sub>9</sub>N<sub>3</sub>S<sub>2</sub>: C, 42.18%; H, 4.55%; N, 21.09%; and Observed values: C, 40.66%; H, 3.49%; N, 22.15%.

The UV spectra of the ligand L<sub>1</sub> shows 3 bands ( $\lambda_{max}$ ) at 210, 256 and 326 nm. The first two bands at 210 and 256 nm are attributed to  $\pi \rightarrow \pi^*$  transitions of the aromatic ring, and thioketonic moieties, respectively<sup>22</sup>. The other band at 326 nm due to  $n \rightarrow \pi^*$  is associated with the azomethine linkage<sup>23,24</sup>. FT-IR spectra for the ligand L<sub>1</sub> is given in Figure 6. Infrared spectra of the ligands show strong bands at 3407 cm<sup>-1</sup> and 3146 cm<sup>-1</sup> were assigned to symmetric or asymmetric  $\nu(\text{NH}_2)$  stretching and  $\nu(\text{N-H})$  vibration of free NH<sub>2</sub> groups<sup>25,26</sup>. The spectrum also shows bands

at 1588, 1294 and 838 cm<sup>-1</sup> due to the  $\nu(\text{C=N})$ ,  $\nu(\text{C=S})$  &  $\delta(\text{C=S})$  groups respectively<sup>27,28</sup>.

The <sup>1</sup>H NMR spectra of the ligand exhibits signals are given as (ppm)  $\delta$ 10.35 (H,s,- NH);  $\delta$ 7.43-7.58 (m,Ar-H); 8.3 (H,s,- N=CH); 3.39 (2H,s,-NH<sub>2</sub>);  $\delta$ 2.28 (s, H<sub>3</sub>C-C)<sup>29</sup>.

**thiophene-2-aldehyde thiosemicarbazone (L<sub>2</sub>)-** Analytical calculation (%) for = C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>S<sub>2</sub>: C, 38.89%; H, 3.81%; N, 22.68%; and Observed values: C, 41.35%; H, 3.68%; N, 24.05%.

The UV spectra of the ligand L<sub>2</sub> shows 3 bands ( $\lambda_{max}$ ) at 208, 262 and 329 nm. The first two bands at 208 and 262 nm are attributed to  $\pi \rightarrow \pi^*$  transitions of the aromatic ring, and thioketonic moieties, respectively<sup>22</sup>. The other band at 329 nm due to  $n \rightarrow \pi^*$  is associated with the azomethine linkage<sup>23,24</sup>. IR spectra for the ligand L<sub>2</sub> is given in Figure 7. Infrared spectra of the ligands show strong bands at 3413 cm<sup>-1</sup> and 3135 cm<sup>-1</sup> were assigned to symmetric or asymmetric  $\nu(\text{NH}_2)$  stretching and  $\nu(\text{N-H})$  vibration of free NH<sub>2</sub> groups<sup>25,26</sup>. The spectrum also shows bands at 1579, 1279 and 837 cm<sup>-1</sup> due to the  $\nu(\text{C=N})$ ,  $\nu(\text{C=S})$  &  $\delta(\text{C=S})$  groups respectively<sup>27,28</sup>.

The <sup>1</sup>H NMR spectra of the ligand exhibits signals are given as (ppm) –  $\delta$ 11.33 (H,s,- NH); 6.94-7.41 (Ar-H); 8.14 (H,s,- N=CH); 2.92 (2H,s,-NH<sub>2</sub>)<sup>29</sup>.

#### Metal Complexes of ligand L<sub>1</sub>

In FT-IR spectra of metal complex (Fig. 8 & 9), the position of ligand band due to azomethine moiety ( $>\text{C=N}$ ) at 1588 cm<sup>-1</sup> shifted towards lower side i.e. 1492 cm<sup>-1</sup> compared to the uncoordinated ligand, indicating its involvement in coordination with metal ion. The ( $>\text{C=S}$ ) stretching frequency observed at 1294 cm<sup>-1</sup> in ligand was shifted in the spectra of the complex at 1328cm<sup>-1</sup>, indicating the involvement of the sulphur in the coordination<sup>24, 28</sup>. This indicates, that the coordination take place through the nitrogen atom of the imine groups and sulphur atom of the thioketonic group ( $>\text{C=S}$ ). In the <sup>1</sup>H NMR spectrum, signals due to -NH- in ligand undergoes downfield shift on complexation with metal. This indicates participation of nitrogen and thioketonic sulphur in bonding.

#### Assessment of Antibacterial assay

All compounds showed significant antibacterial activity at the concentration of 50 µg/ml (Table 1) assured by Kirby-Bauer's method. The inhibition potential of bacterial growth exhibited by all compounds has shown in Figure 4 & 5, observation revealed that synthesized lead complex of L<sub>1</sub> ligand, ([Pb(L<sub>1</sub>)<sub>2</sub>]Cl<sub>2</sub>) has much stronger anti-bacterial effect against all compounds. Thus, the synthesized metal complex of ligand L<sub>1</sub> acts as excellent antibacterial activity against both gram positive and gram-negative bacteria with zone of inhibition 12.45 ± 1.2 and 11.27 ± 1.1 mm respectively.


## CONCLUSION

In the present work we successfully synthesized 2-acetylthiophene thiosemicarbazone and thiophene-2-aldehyde thiosemicarbazone ligands and their metal complexes. These ligands and metal complexes have been characterized by various analytical and spectral techniques. The results of antibacterial activity reveals that these compounds exhibited significant activity against *E. Coli* and gram-positive bacterial strains *Bacillus sp*.


**Table 1: Antibacterial activity (% inhibition) of ligands complexes against *E. coli*. and *Bacillus sp* at various concentrations**

S. No.	Zone of Inhibition (Diameter in mm)			
	Treatment	Concentration (in µg)	<i>Escherichia coli</i>	<i>Bacillus sp</i>
1	Normal control	-	1.2 ± 0.2	1.1 ± 0.2
2	Ampicillin	10	14.55 ± 0.9 <sup>a</sup>	15.23 ± 0.7 <sup>a</sup>
3	L <sub>1</sub>	50	9.67 ± 0.8 <sup>a</sup>	11.33 ± 0.8 <sup>a</sup>
4	L <sub>2</sub>	50	9.3 ± 0.8 <sup>a</sup>	9.6 ± 0.8 <sup>a</sup>
5	[(Pb(L <sub>1</sub> ) <sub>2</sub> )Cl <sub>2</sub> ]	50	12.45 ± 1.2 <sup>a</sup>	11.27 ± 1.1 <sup>a</sup>
6	[Ni(L <sub>2</sub> ) <sub>2</sub> ]Cl <sub>2</sub>	50	9.1 ± 0.9 <sup>a</sup>	11.2 ± 0.8 <sup>a</sup>


All data has represented in mean ± SD. a = *p* < 0.05 vs normal control, b = *p* < 0.05 vs standard concentration (10 µg) of ampicillin in respective bacterial strain




**Figure 1: Scheme for the synthesis of 2-acetylthiophene thiosemicarbazone (L1)**



**Figure 2: Scheme for the synthesis of thiophene-2-aldehyde thiosemicarbazone (L2)**



**Figure 3: Synthesis of metal complexes [M(L)<sub>2</sub>]Cl<sub>2</sub>**



**Figure 4: Zone of inhabitation against *Escherichia coli***






Figure 9: FT-IR and  $^1\text{H}$  NMR spectrum of Nickel(II) complex with ligand L1

## ACKNOWLEDGEMENTS

The authors are thankful to the University Grant Commission, New Delhi, for providing financial assistance and the Management, K. R. Mangalam University for providing research facilities. We also express our sincere thanks to IIT Delhi, for recording FT-IR and NMR spectra.

## REFERENCES

- Dinku W, Megersa N, Raju VJT, Solomon T, Jonsson JA and Retta N. Studies of transition metal complexes of herbicidal compounds. I: Transition metal complexes of 2-methylthio-4-azido-6-isopropylamino-s-triazine (aziprotryn). *Bulletin of the Chemical Society of Ethiopia* 2001; 15(1): 29-37. <https://doi.org/10.4314/bcse.v15i1.71731>
- Gaur S. Physico-Chemical and Biological Properties of Mn(II), Co(II), Ni(II) and Cu(II) Chelates of Schiff Bases. *Asian Journal of Chemistry* 2003; 15 (1): 250-254. (b) Genin MJ, Biles C, Keiser BJ, Poppe SM, Swaney SM, Tarapley WG et. al. Novel 1,5 Diphenylpyrazole Nonnucleoside HIV-1 Reverse Transcriptase Inhibitors with Enhanced Activity versus the Delavirdine-Resistant P236L Mutant: Lead Identification and SAR of 3- and 4-Substituted Derivatives. *Journal of Medicinal Chemistry* 2000; 43 (5): 1034-1040. <https://doi.org/10.1021/jm990383f>
- Rao R, Patra AK, Chetana PR. DNA binding and oxidative cleavage activity of ternary (L-proline) copper (II) complexes of heterocyclic bases. *Polyhedron* 2007; 26: 5331-5338. <https://doi.org/10.1016/j.poly.2007.07.040>
- Karon M, Benedict WF. Chromatid breakage: differential effect of inhibitors of DNA synthesis during G2 phase. *Science* 1972; 178: 69. <https://doi.org/10.1126/science.178.4056.62>
- Jungreis E, Thabet S. *Analytical Applications of Schiff bases*. Marcell Dekker: New York; 1969. PMID:5357675
- Sari N, Arslan S, Logoglis E, Sakiyan. Synthesis, Characterization, and Antibacterial Activity of the Schiff Bases Derived from Thiosemicarbazide, Salicylaldehyde, 5-bromosalicylaldehyde and their Copper(II) and Nickel(II) Complexes. *IGU Journal of Science* 2003; 283.
- Amir M, Hasan SM, Wadood A. Synthesis and Antibacterial Activity of 1-Isonicotinyl-3-Methyl-4-(Substituted Phenyl Hydrazono)-2-Pyrazolin-5-one. *Oriental Journal of Chemistry* 2002; 18 (2): 351-355.
- Sridhar SK, Saravaran M, Rameshver. Synthesis and antibacterial screening of hydrazones, Schiff and Mannich bases of isatin derivatives. *Eurasian Journal of Medicinal Chemistry* 2001; 36 (7-8): 615-625. [https://doi.org/10.1016/S0223-5234\(01\)01255-7](https://doi.org/10.1016/S0223-5234(01)01255-7)
- Rajendran SP, Karvembu R. Synthesis and antifungal activities of Schiff bases derived from 3-amino-2H-pyran [2,3-b]-quinolin-2-ones. *Indian Journal of Chemistry* 2002; 41B: 222. <https://doi.org/10.1002/chin.200219139>
- Calis U, Yarim M, Kokasal M, Ozalp M. Synthesis and antimicrobial activity evaluation of some new adamantane derivatives. *Arzneimittel-Forschung* 2002; 52(10): 778-781.
- Pandeya SN, Sriram D, Nath G. Synthesis and antimicrobial activity of Schiff and Mannich bases of isatin and its derivatives with pyrimidine. *Farmaco* 1999; 54: 624- 628. [https://doi.org/10.1016/S0014-827X\(99\)00075-0](https://doi.org/10.1016/S0014-827X(99)00075-0)
- Shaikh KA, Baseer MA, Mote NA. *Asian Journal of Chemistry* 2001; 13: 496.
- Karia FD, Parsania PH. Synthesis, biological and thermal properties of Schiff bases of bisphenol, C. *Asian Journal of Chemistry* 1999; 11(3): 991-995.
- Sridhar SK, Pandaya SN, De Cleroq E. *Bollettino Chimico Farmaceutico* 2001; 140: 302. PMID:11680082
- Isaacs NS. *Chemical Society Reviews* 1976; 5: 181. <https://doi.org/10.1039/CS9760500181>
- Schmid GH. *Organic Chemistry*. Mosby New York; 1996: 624.
- Mohan C, Sharma K and Kumari S. Synthesis and Structural Studies of Schiff based Metal Complexes of Semicarbazones and Thiosemicarbazones. *International Journal of Chemical and Natural Science* 2015; 3 (5): 312-319.
- Mary NL and Parameswaran G. Synthesis and Reactivity in *Inorganic and Metal-Organic Chemistry* 1993; 23(7): 1209–1227. <https://doi.org/10.1080/15533179308016680>
- Mary NL and Parameswaran G. *Croatica Chemica Acta* 1993; 66(2): 355–362.
- Kumar V, Sharma A, Rajput S, Pal M and Dhiman N. Pharmacognostic and pharmacological evaluation of *Eulaliopsis binata* plant extracts by measuring in vitro/in vivo safety profile and anti-microbial potential. *Toxicological Research* 2018; 7: 454-464. <https://doi.org/10.1039/C8TX00017D>
- Mohan C, Sharma K and Chandra S. Cd (II) ion-selective electrode based on 2-acetylthiophene Semicarbazone in Polymeric membrane. *Analytical & Bioanalytical Electrochemistry* 2017; 9 (1): 35-46.
- Eisenmann G. *The electrochemistry of cation selective glass electrodes in advances in analytical chemistry and instrumentation*. John Wiley and sons. Inc: New York; 1965.
- Eisenmann G. *Glass electrodes for hydrogen and other cations*. Principles and practice. Dekker M. New York; 1967.
- Tendeloo HJC. A new and easy method for the potentiometric determination of calcium concentrations in solutions. *Journal of Biological Chemistry* 1936; 113(333): 197.
- Ramosa JM, Viana RM, Tellez CA et al. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 2006; 65: 433–438. <https://doi.org/10.1016/j.saa.2005.09.047> PMID:16931127

26. Maurya RC and Mishra DD. Synthesis and characterization of some novel mixed-ligand cyanonitrosyl  $\{CrNO\}_5$  heterocomplexes of chromium(I) with potentially tridentate benzothiazole derivatives. *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry* 1992; 22: 1227–1237. <https://doi.org/10.1080/15533179208020264>
27. Chandra S and Gupta LK. Spectroscopic approach in characterization of chromium(III), manganese(II), iron(III) and copper(II) complexes with a nitrogen donor tetradentate, 14-membered azamacrocyclic ligand. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 2005; 61(9): 2139-2144. <https://doi.org/10.1016/j.saa.2004.06.060> PMID:15911403
28. Ali MA, Mirza AH, Ejau WB and Bernhardt PV. Preparation and characterization of pentagonal-bipyramidal zinc (II) and cadmium (II) complexes of a neutral N<sub>3</sub>S<sub>2</sub> thiosemicarbazone derived from 2 N-methylthiocynate. *Polyhedron* 2000; 25 (17): 3337-3342.
29. Mundwiler S, Candrea L, Häfliger P, Ortner K and Alberto R. Preparation of NO-Carrier-Added Technetium-99m Complexes via Metal-Assisted Cleavage from a Solid Phase. *Bioconjugate Chemistry* 2004; 15: 195–202. <https://doi.org/10.1021/bc034171f> PMID:14733600
30. Gambino D, Otero L, Vieitesa M, Boiani M et al. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy* 2007; 68: 341–348. <https://doi.org/10.1016/j.saa.2006.11.043> PMID:17321785

**Cite this article as:**

Chandra Mohan *et al.* Synthesis, characterization, and antibacterial activity of Schiff bases derived from thiosemicarbazide, 2-acetyl thiophene and thiophene-2 aldehyde. *Int. Res. J. Pharm.* 2018;9(7):153-158 <http://dx.doi.org/10.7897/2230-8407.097141>

Source of support: University Grant Commission, New Delhi, India, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.