



## Research Article

## AN ALTERNATIVE TOTAL SYNTHESIS OF (–)-PYRENOPHOROL

Sridhar Musulla<sup>1,2</sup>, Bharathi Kumari Y<sup>\*2</sup>, Srinivasa Rao. A<sup>1</sup><sup>1</sup>GVK Biosciences Private Limited, Nacharam, IDA Mallapur, Hyderabad, Telangana, India<sup>2</sup>Department of Chemistry, JNT University, Hyderabad, Telangana, India

\*Corresponding Author Email: sridharmusulla@rediffmail.com

Article Received on: 06/08/18 Approved for publication: 12/09/18

DOI: 10.7897/2230-8407.099206

## ABSTRACT

Aim: The aim of the present work is to report an alternative strategy to synthesis of (–)-Pyrenophorol. Methods: In this synthesis all intermediates and target compounds were fully characterized by different spectroscopic techniques such as <sup>1</sup>H NMR, <sup>13</sup>CNMR and EI-MS. Results and Discussion: The total synthesis of 16-membered C<sub>2</sub>-symmetric dilactone (–)-Pyrenophorol was prepared from (R, S)-di epoxide in 9 steps with good yields using Wittig olefination and Mitsunobu cyclization as key steps. Conclusion: A short and efficient stereoselective total synthesis of Pyrenophorol (1) has been achieved from the known di epoxide. The key steps includes Wittig olefination and dimerization by Mitsunobu cyclization.

**Keywords:** pyrenophorol, Ozonolysis, Wittig olefination, cyclodimerization, Mitsunobu cyclization


## INTRODUCTION

Pyrenophorol (1), which belongs to the family of symmetrical macrodiolides, isolated from the fungus *Byssoschlamys nivea*<sup>1</sup>, and *Stemphylium radicinum*<sup>2</sup>. It was later isolated from *Alternaria alternata*<sup>3</sup>, *Drechslera avenae*<sup>4</sup> and *Phoma* sp.<sup>5</sup>. Diolide 1 exhibits pronounced anthelmintic properties<sup>3</sup>, besides moderate activity against the fungus *Microbotryum violaceum*.

Pyrenophorol (1) is a 16-membered bis-lactone embedded with four stereocenters (5S, 8R, 13S, 16R) and two conjugated double bonds. Structurally 1 is related to pyrenophorin<sup>6</sup>, and vermiculin<sup>7</sup>, while, the monomeric unit,  $\gamma$ -hydroxy- $\alpha,\beta$ -unsaturated ester is structurally related to 14-membered macrodiolide colletello<sup>8</sup>, Zwanenburg *et al.*<sup>9</sup> reported the synthesis of 5R, 8R, 13R, 16R and 5S, 8R, 13S, 16R isomers of diolide and determined the absolute stereochemistry of 1 as 5S, 8R, 13S, 16R based on the spectral, analytical and optical rotation data.

The structural features coupled with interesting biological activities, prompted several synthesis for 1 by different synthetic strategies<sup>9, 10</sup> while, Floch and Amigoni<sup>11</sup> reported the synthesis of 5R, 8S, 13R, 16S stereoisomer. Zwanenburg *et al.*<sup>9</sup> and Kang *et al.*<sup>12</sup> utilized Yamaguchi method<sup>14</sup> for macrolide synthesis, while, Kibayashi *et al.*<sup>10</sup> and Yadav *et al.*<sup>11</sup> adopted dimerization of hydroxy acid under Mitsunobu conditions<sup>15</sup>, whereas, Floch, and Amigoni<sup>13</sup> method of macrocyclization was achieved through intramolecular Wittig reaction.

In continuation of our interest on the synthesis of macrodiolides<sup>16-19</sup> herein, we report the synthesis of pyrenophorol (1) by adopting dimerization of hydroxy acid using Mitsunobu protocol.



Pyrenophorol (1)  
(5S, 8R, 13S, 16R)

Figure 1

## MATERIALS AND METHODS

Solvents were dried over standard drying agents on freshly distilled prior to use. Chemicals were purchased and used without further purification. All column chromatographic separations were performed using silica gel (60-120 mesh). Organic solutions were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated below 40 °C in *vacuo*. <sup>1</sup>H NMR spectra were acquired at 300 MHz, 500 MHz and 600 MHz, while, <sup>13</sup>C NMR at 75 MHz and 125 MHz with TMS as internal standard for solutions in CDCl<sub>3</sub>. *J* values were given in Hz. IR-spectra were recorded on FT IR spectrophotometer with NaCl optics. Optical rotations were measured on digital polarimeter at 25 °C. Mass spectra were recorded on direct inlet system or LC by MSD trap SL, the HRMS data were obtained using Q-TOF mass spectrometry.

(3S,6S)-6-(*tert*-Butyldimethylsilyloxy)hept-1-en-3-ol (3)

To a suspension of trimethylsulfonium iodide (5.70 g, 27.7 mmol) in dry THF (20 mL) at -10 °C under nitrogen atmosphere was added *n*-BuLi (2.5M in hexane, 9.3 mL, 23.3 mmol). After 30 min, compound 6 (3.0 g, 9.3 mmol) in THF (10 mL) was

introduced, producing a milky suspension. The reaction was allowed to warm to 0 °C over 30 minutes and then to room temperature and stirred for 4 hours. The reaction was quenched with water (20 mL) at 0 °C, extracted with EtOAc (2 x 50 mL) and the combined organic layers dried over sodium sulfate. The crude residue was purified by column chromatography (60-120 mesh Silica gel) to afford the secondary alcohol **3** (2.2 g) in 70% yield.  $[\alpha]_D^{25}$  -37.4 (*c* 0.18, CHCl<sub>3</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 5.81 (m, 1H, olefinic), 5.13 (d, 1H, *J* = 14.8 Hz, olefinic), 5.01 (d, 1H, *J* = 10.4 Hz, olefinic), 3.99 (m, 1H, -CH), 3.78 (m, 1H, -CH), 1.60-1.37 (m, 4H, 2 x -CH<sub>2</sub>), 1.06 (d, 3H, *J* = 5.4 Hz, -CH<sub>3</sub>), 0.81 (s, 9H, 3 x -CH<sub>3</sub>), 0.01 (s, 6H, 2 x -CH<sub>3</sub>); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 141.5, 114.3, 72.1, 68.6, 33.1, 26.0, 23.3, 18.0, -4.8, -4.4; IR (neat): 3435, 2929, 2857, 1465, 1373, 1253, 1134, 1048, 833 cm<sup>-1</sup>; ESIMS *m/z* (M+Na)<sup>+</sup>: 267

**tert.-Butyl((2S,5S)-5-(4-methoxybenzyloxy)hept-6-en-2-yloxy)dimethylsilane (7)**

To a cooled (0 °C) solution of **3** (3.0 g, 12.29 mmol) in dry THF (30 mL), NaH (0.59 g, 24.59 mmol) was added, stirred for 30 min and treated with a solution of PMBBr (2.93 g, 14.74 mmol) in dry THF (15 mL). After stirring 7.5 h at room temperature, it was quenched with sat. NH<sub>4</sub>Cl solution (10 mL) and extracted with ethyl acetate (2 x 50 mL). Organic layers were washed with water (2 x 10 mL), brine (10 mL) and dried (Na<sub>2</sub>SO<sub>4</sub>). Solvent was evaporated under reduced pressure and purified the residue by column chromatography (60-120 Silica gel, 5% EtOAc in pet. ether) to furnish **7** (3.7 g, 82%) as a yellow liquid.  $[\alpha]_D^{25}$  -23.6 (*c* 0.9, CHCl<sub>3</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.20 (d, 2H, *J* = 8.6 Hz, ArH-PMB), 6.83 (d, 2H, *J* = 8.6 Hz, ArH-PMB), 5.77-5.61 (heptet, 1H, *J* = 7.5, 10.3 Hz, olefinic), 5.19 (q, 2H, *J* = 4.1, 10.3 Hz, olefinic), 4.54, 4.30 (2d, 2H, *J* = 11.8 Hz, -OCH<sub>2</sub> Ar), 3.78 (s, 3H, -OCH<sub>3</sub>), 3.76-3.62 (m, 2H, 2 x -CH), 1.61-1.32 (m, 4H, 2 x -CH<sub>2</sub>), 1.20 (d, 3H, *J* = 6.0 Hz, -CH<sub>3</sub>), 0.85 (s, 9H, 3 x -CH<sub>3</sub>) 0.00 (s, 6H, 2 x -CH<sub>3</sub>); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 149.0, 131.5, 128.5, 128.2, 127.6, 121.0, 72.7, 57.80, 55.60, 35.3, 30.2, 25.8, 23.8, 22.4, -4.4; IR (neat): 2926, 2858, 1722, 1456, 1268, 1106 cm<sup>-1</sup>; ESIMS *m/z* (M+Na)<sup>+</sup>: 387

**(4S,7S,E)-Methyl 7-(tert.-butyldimethylsilyloxy)-4-(4-methoxybenzyloxy)oct-2-enoate (8)**

Ozone was bubbled through a cooled (-78 °C) solution of **7** (7.4 g, 34.57 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (70 mL) until the pale blue color persisted. Excess ozone was removed with Me<sub>2</sub>S (2 mL) and stirred for 15 min at 0 °C. The reaction mixture was concentrated under reduced pressure to give aldehyde, which was used for further reaction.

A solution of the above aldehyde in benzene (50 mL) was treated with (methoxy- carbonylmethylene)triphenyl phosphorane (3.54 g, 10.54 mmol) at reflux temperature. After 2 h, solvent was evaporated and purified the residue by column chromatography (60-120 Silica gel, 10% EtOAc in pet. ether) to furnish **8** (3.12 g, 84%) as a yellow liquid.  $[\alpha]_D^{25}$  -48.6 (*c* 1.0, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.16 (d, 2H, *J* = 8.3 Hz, ArH-PMB), 6.81 (d, 2H, *J* = 8.1 Hz, ArH-PMB), 6.77 (dd, 1H, *J* = 6.1, 15.8 Hz, olefinic), 5.94 (d, 1H, *J* = 15.6 Hz, olefinic), 4.47 (d, 1H, *J* = 11.7 Hz, benzylic), 4.24 (d, 1H, *J* = 11.7 Hz, benzylic), 3.87 (q, 1H, *J* = 5.6, 12.1 Hz, -OCH), 3.77 (s, 3H, OCH<sub>3</sub>), 3.73 (s, 3H, OCH<sub>3</sub>), 3.61 (m, 1H, -OCH), 1.72-1.31 (br. m, 4H, 2 x -CH<sub>2</sub>), 1.07 (d, 3H, *J* = 6.0 Hz, -CH<sub>3</sub>), 0.85 (s, 9H, 3 x -CH<sub>3</sub>), 0.01 (s, 6H, 2 x -CH<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 150 MHz): δ 166.5, 158.2, 147.4, 129.7, 128.6, 118.8, 113.5, 79.6, 71.2, 66.6, 55.4, 51.6, 35.4, 30.2, 25.6, 24.2, 18.2, -4.8; IR (neat): 2932, 1724, 1612, 1512, 1448, 1386, 1164, 1037 cm<sup>-1</sup>; ESIMS *m/z* (M+Na)<sup>+</sup>: 445

**(4S,7S,E)-7-(tert.-Butyldimethylsilyloxy)-4-(4-methoxybenzyloxy)oct-2-enoic acid (9)**

To a solution of **8** (2.6 g, 6.16 mmol) in THF: MeOH: water (3:1:1, 20 mL), LiOH (0.45 g, 18.48 mmol) was added and stirred at room temperature for 4 h. The pH of reaction mixture was adjusted to acidic with 1N HCl solution and extracted with ethyl acetate (30 mL). Organic layers were washed with water (15 mL), brine (15 mL) and dried (Na<sub>2</sub>SO<sub>4</sub>). Solvent was evaporated under reduced pressure and purified the residue by column chromatography (60-120 Silica gel, 30% EtOAc in pet. ether) to afford **9** (2.02 g, 80%) as a colourless oil.  $[\alpha]_D^{25}$  +14.6 (*c* 0.6, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.19 (d, 2H, *J* = 8.5 Hz, ArH-PMB), 6.94 (dd, 1H, *J* = 6.0, 15.6 Hz, olefinic), 6.81 (d, 2H, *J* = 8.5 Hz, ArH-PMB), 5.91 (d, 1H, *J* = 15.5 Hz, olefinic), 4.48 (d, 1H, *J* = 11.4 Hz, benzylic), 4.29 (d, 1H, *J* = 11.6 Hz, benzylic), 3.90 (q, 1H, *J* = 5.6, 12.1 Hz, -OCH), 3.79 (s, 3H, OCH<sub>3</sub>), 3.49 (m, 1H, -OCH), 1.72-1.34 (br. m, 4H, 2 x -CH<sub>2</sub>), 1.13 (d, 3H, *J* = 6.0 Hz, -CH<sub>3</sub>); 0.85 (s, 9H, 3 x -CH<sub>3</sub>), 0.09 (s, 6H, 2 x -CH<sub>3</sub>); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ 172.8, 158.4, 149.5, 130.6, 128.6, 119.6, 113.5, 78.8, 72.6, 66.6, 55.6, 36.2, 30.8, 25.9, 24.2, 17.6, -4.6; IR (neat): 3540, 3031, 2930, 2857, 1710, 1097 cm<sup>-1</sup>; ESIMS *m/z* (M+Na)<sup>+</sup>: 431

**(4S,7S,E)-7-Hydroxy-4-(4-methoxybenzyloxy)oct-2-enoic acid (2)**

To a cooled (0 °C) solution of **9** (2.20 g, 5.40 mmol) in dry THF (15 mL) under nitrogen atmosphere, TBAF (6.5 mL, 6.5 mmol) was added and stirred for 3 h. Reaction mixture was diluted with water (5 mL) and extracted with ethyl acetate (2 x 50 mL). Organic layers were washed with water (2 x 10 mL), brine (10 mL) and dried (Na<sub>2</sub>SO<sub>4</sub>). Solvent was evaporated and purified the residue by column chromatography (60-120 Silica gel, 55% EtOAc in pet. ether) to furnish **2** (1.38 g, 87%) as a liquid.  $[\alpha]_D^{25}$  -32.6 (*c* 1.0, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.18 (d, 2H, *J* = 8.6 Hz, ArH-PMB), 6.95 (dd, 1H, *J* = 6.0, 15.6 Hz, olefinic), 6.80 (d, 2H, *J* = 8.6 Hz, ArH-PMB), 5.99 (d, 1H, *J* = 15.6 Hz, olefinic), 4.50 (d, 1H, *J* = 11.6 Hz, benzylic), 4.29 (d, 1H, *J* = 11.6 Hz, benzylic), 3.92 (q, 1H, *J* = 5.6, 12.1 Hz, -OCH), 3.79 (s, 3H, OCH<sub>3</sub>), 3.76 (m, 1H, -OCH), 1.72-1.32 (br. m, 4H, 2 x -CH<sub>2</sub>), 1.11 (d, 3H, *J* = 6.0 Hz, -CH<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz): δ 172.6, 158.2, 146.4, 132.6, 128.9, 118.3, 113.6, 78.8, 72.4, 68.2, 56.4, 34.2, 29.8, 23.2; IR (neat): 3451, 2929, 2857, 2102, 1722, 1612, 1514, 1360, 1041, 777 cm<sup>-1</sup>; ESIMS *m/z* (M+Na)<sup>+</sup>: 317.1355.

**(3E,5S,8R,11E,13S,16R)-5,13-Bis(4-methoxybenzyloxy)-8,16-dimethyl-1,9-dioxacyclo-hexadeca-3,11-diene-2,10-dione (10)**

A solution of **2** (0.225 g, 0.93 mmol) and Ph<sub>3</sub>P (1.22 g, 4.67 mmol) in toluene: THF (10:1, 250 mL), DEAD (0.81 mL, 16.87 mmol) was added at -20 °C and stirred under N<sub>2</sub> atmosphere for 10 h. Solvent was evaporated under reduced pressure and purified the residue by column chromatography (60-120 Silica gel, 10% EtOAc in pet. ether) to afford **10** (0.14 g, 56%) as a colorless oil.  $[\alpha]_D^{25}$  -15.7 (*c* 1.03, CHCl<sub>3</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): 7.18 (d, 4H, *J* = 8.7 Hz, ArH-PMB), 6.81 (d, 4H, *J* = 8.7 Hz, ArH-PMB), 6.61 (dd, 2H, *J* = 6.2, 11.6 Hz, olefinic), 5.82 (d, 2H, *J* = 11.6 Hz, olefinic), 5.01-4.91 (m, 2H, -OCH), 4.41 (d, 2H, *J* = 11.3 Hz, benzylic), 4.24 (d, 2H, *J* = 11.3 Hz, benzylic), 4.12 (m, 2H, -OCH), 3.61 (s, 6H, OCH<sub>3</sub>), 1.79 (q, 4H, *J* = 6.4 Hz, -CH<sub>2</sub>), 1.60 (m, 4H, -CH<sub>2</sub>), 1.32 (d, 6H, *J* = 7.1 Hz, -CH<sub>3</sub>); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): 166.6, 158.2, 146.2, 129.7, 128.8, 120.6, 113.6, 79.8, 72.4, 68.4, 56.2, 39.6, 28.4, 2.2; IR (neat): 3416, 3068, 2932, 2859, 1722, 1608, 1527, 1462, 1427, 1273, 1105, 918, 702 cm<sup>-1</sup>; ESIMS *m/z* (M+Na)<sup>+</sup>: 575


**Pyrenophorol (1)**

To a solution of **10** (0.12 g, 0.21 mmol) in aq.  $\text{CH}_2\text{Cl}_2$  (2 mL, 19:1), DDQ (71 mg, 0.31 mmol) was added and stirred at room temperature for 3 h. The reaction mixture was quenched with sat.  $\text{NaHCO}_3$  solution (1 mL), filtered and washed with  $\text{CH}_2\text{Cl}_2$  (10 mL). The filtrate was washed with water (3 mL), brine (3 mL) and dried ( $\text{Na}_2\text{SO}_4$ ). Solvent was evaporated under reduced pressure and purified the residue by column chromatography (60-120 Silica gel, 20% EtOAc in pet. ether) to furnish **1** (52 mg, 78%) as a white solid. m.p.: 137-139 °C; lit.<sup>1</sup> m.p. 135 °C;  $[\alpha]_{\text{D}} -3.7$  (*c* 0.13, acetone); lit.<sup>1</sup>  $[\alpha]_{\text{D}}^{25} -3.0$  (*c* 1.0, acetone);  $^1\text{H NMR}$  ( $\text{CDCl}_3$ , 300 MHz):  $\delta$  6.81 (dd, 2H,  $J = 15.4, 5.1$  Hz, olefinic), 5.87 (dd, 2H,  $J = 15.4, 2.0$  Hz, olefinic), 5.11-4.99 (m, 2H, 2 x -OCH), 4.21-4.19 (m, 2H, 2 x -OCH), 2.29 (br. s, 2H, 2 x -OH),

2.03-1.93 (m, 4H, 2 x - $\text{CH}_2$ ), 1.77-1.58 (m, 4H, 2 x - $\text{CH}_2$ ), 1.23 (d, 6H,  $J = 6.0$  Hz, 2 x - $\text{CH}_3$ );  $^{13}\text{C NMR}$  (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  167.5, 143.3, 121.3, 74.1, 68.6, 31.1, 28.9, 19.0; IR (neat): 3442, 2922, 2853, 1721, 1630, 1126, 835  $\text{cm}^{-1}$ ; ESIMS  $m/z$  ( $\text{M}+\text{Na}^+$ ): 335

**RESULTS AND DISCUSSION**


Retrosynthetic analysis of macrodiolide **1** reveal that they could be obtained from the hydroxy acids **2** via cyclodimerisation under the Mitsunobu conditions followed by deprotection of PMB ethers. Hydroxy acid **2** could be prepared from the racemic allylic alcohol **3**, which could be prepared from di epoxide **4** by simple chemical transformations.



Scheme 1

The synthesis of macrodiolide **1** was initiated from (*R,S*)-di epoxide<sup>20</sup> as illustrated in Scheme 2. Accordingly, the di epoxide **4** was opened regioselectively with DIBAL-H in  $\text{CH}_2\text{Cl}_2$  at 0 °C to rt and subsequent silylation of the resulting secondary alcohol **5** with TBSCl and imidazole in  $\text{CH}_2\text{Cl}_2$  gave **6** in 70% yield. Later, the second epoxide ring in compound **6** was also opened

with trimethylsulfonium iodide in the presence of *n*-BuLi in THF at -0 °C for 6 h to give the alcohol **3**, which on treatment with NaH and *p*-methoxybenzyl bromide at 0 °C furnished the PMB ether **7** in 82% yield. Next, Ozonolysis of **7** followed by Wittig olefination of the resulting aldehyde afforded **8** in 76% yield.



Scheme 2

**Reagents and conditions:** (a) DIBAL-H, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to rt, 2 h; (b) TBSCl, Imidazole, CH<sub>2</sub>Cl<sub>2</sub>, rt, 4 h; (c) *n*-BuLi, Me<sub>3</sub>Si, THF, 0 °C, 6 h (d) PMBBBr, NaH, THF, 0 °C to rt, 8 h; (e) i) O<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C, 30 min; ii) Ph<sub>3</sub>P=CHCOOMe, Benzene, reflux, 2 h; (f) LiOH, THF:MeOH:H<sub>2</sub>O (3:1:1), rt, 4 h; (g) TBAF, THF, 0 °C to rt, 3 h; (h) Ph<sub>3</sub>P, DEAD, toluene:THF (10:1) -25 °C, 10 h; (i) DDQ, CH<sub>2</sub>Cl<sub>2</sub>:H<sub>2</sub>O (19:1), rt, 3 h.

Ester **8** on subsequent hydrolysis (LiOH in THF:MeOH:H<sub>2</sub>O-3:1:1) gave acid **9**, which on desilylation with TBAF in dry THF afforded the hydroxy-acid **2** in 87% yield. Hydroxy-acid **2** on cyclodimerization under Mitsunobu reaction conditions according to Gerlach's procedure [13] with Ph<sub>3</sub>P and DEAD at -25 °C for 10 h furnished **10**. Finally, oxidative deprotection of PMB groups in **10** using DDQ in aq. CH<sub>2</sub>Cl<sub>2</sub>:H<sub>2</sub>O (19:1) provided (-)-pyrenophorol (**1**) in 78% yield. [α]<sub>D</sub> -3.8 (c 0.66, acetone); lit.<sup>1,9</sup> [α]<sub>D</sub> -3.0 (c 1.0, acetone); The <sup>1</sup>H and <sup>13</sup>C NMR data and optical rotation value of synthetic **1** is identical to those reported in the literature.<sup>10</sup>

## CONCLUSION

Thus, in summary a short and efficient stereoselective total synthesis of Pyrenophorol (**1**) has been achieved from the known di epoxide. The key steps includes Wittig olefination and dimerization by Mitsunobu cyclization.

## ACKNOWLEDGEMENT

The author is thankful to JNT University for constant encouragement during this research program. The author is also thankful to GVK Biosciences for providing necessary facility for research work.

## REFERENCES

- Kis Z, Furger P, Sigg H. Über die Isolierung von Pyrenophorol. *Experientia* 1969; 25: 123-124.
- Grove FJF. Metabolic products of *Stemphylium radicinum*. Part IV. Minor products. *Journal of the Chemical Society C* 1971; 2261-2263.
- Kind R, Zeeck A, Grabley S, Thiericke R, Zerlin MJ. Secondary Metabolites by Chemical Screening. Helmidol, a New Macrodiolide from *Alternaria alternate*. *Journal of Natural Products* 1996; 59:539-540.
- Kastanias MA, Chrysayi-Tokousbalides M. Herbicidal potential of pyrenophorol isolated from a *Drechslera avenae* pathotype. *Pest Management Science* 2000; 56: 227-232.
- Krohn K, Farooq U, Flörke U, Schulz B, Draeger S, Pescitelli G, Salvadori P, Antus S, Kurtán T. Secondary Metabolites Isolated from an Endophytic *Phoma* sp. – Absolute Configuration of Tetrahydropyrenophorol Using the Solid-State TDDFT CD Methodology. *European Journal of Organic Chemistry* 2007; 3206-3211.
- Nozoe S, Hira K, Tsuda k, ishikashi k, Grove JF. The structure of pyrenophorin. *Tetrahedron Letters* 1965; 4675-4677.
- Noda A, Aoyagi S, Machinaga N, Kibayashi C. Total synthesis of (-)-vermiculine. *Tetrahedron Letters* 1994; 35: 8237-8240.
- MacMillan J, Simpson TJ. Fungal products. Part V. The absolute stereochemistry of colletodiol and the structures of related metabolites of *Colletotrichum capsici*. *Journal of the Chemical Society Perkin I* 1973; 1487-1493.
- Zwanenburg B, Thijs L. Total synthesis of the macrodiolide pyrenophorol. *Tetrahedron Letters* 1991; 32: 1499-1502.
- (a) Kibayashi C, Machinaga N. Preparation of macrodiolides via a common chiral building block. Total synthesis of (-)-pyrenophorin and (-)-pyrenophorol. *Tetrahedron Letters* 1993; 34:841-844.
- Yadav JS, Subba Reddy UV, Subba Reddy BV. Stereoselective total synthesis of (-)-pyrenophorol. *Tetrahedron Letters* 2009; 50: 5984-5986.
- Oh HS, Kang HY. *Bull. Korean Chem. Soc.* 2011; 32:2869.
- Le Floch Y, Amigoni S. A total synthesis of the (5*R*,8*S*,13*R*,16*S*)-isomer of pyrenophorol. *Tetrahedron: Asymmetry* 1997; 8: 2827.
- Inanaga J, Hirata K, Saeki H, Katsuki T, Yamaguchi M. A Rapid Esterification by Means of Mixed Anhydride and Its Application to Large-ring Lactonization. *Bulletin of the Chemical Society of Japan* 1979; 52; 1989.
- Gerlach H, Gertle K, Thahnann A. Eine neue Synthese von (±)-Pyrenophorin. *Helvetica Chimica Acta* 1977; 60: 2860.
- Rao AVR, Murthy V S, Sharma GVM. Studies directed towards the synthesis of clonostachydiol. *Tetrahedron Letters* 1995; 36:139-142.
- Rao AVR, Murthy VS, Sharma GVM. The first synthesis and determination of absolute stereochemistry of clonostachydiol. *Tetrahedron Letters* 1995; 36:143-146.
- Sharma GVM, Goverdhan Reddy C. A stereoselective synthesis of verbalactone—determination of absolute stereochemistry. *Tetrahedron Letters* 2004; 45:7483-7485.
- Sharma GVM, Mallesham S, Chandramouli Ch. Studies directed toward the first total synthesis of acremodiol and acremonol. *Tetrahedron: Asymmetry* 2009; 20:2513-2529.
- Lee SH., Kohn H. Efficient Synthesis of Medium-Sized Cyclic Ether Diamines. *Journal of Organic Chemistry* 2002; 67:1692-1695.

## Cite this article as:

Sridhar Musulla et al. An alternative total synthesis of (-)-Pyrenophorol. *Int. Res. J. Pharm.* 2018;9(9):161-164 <http://dx.doi.org/10.7897/2230-8407.099206>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.