

MEDICINAL PLANTS OF RAJASTHAN (INDIA) WITH ANTIDIABETIC POTENTIAL

Batra Shikha*, Nagori Badri Prakash, Batra Nikhil

Lachoo Memorial College of Science & Technology, Jodhpur, India

Article Received on: 10/01/2011 Revised on: 12/02/2011 Approved for publication: 23/02/2011

*Shikha Batra, Lecturer, Pharmacy Wing, Lachoo Memorial College of Science & Technology, Sector 'A', Shastri Nagar, Jodhpur- 342 003 E mail: shikhabatrapc@yahoo.co.in

ABSTRACT

Rajasthan has a rich heritage of traditional system of medicine and many medicinally useful plants are found growing wildly because of vast area and variety of agro-climatic conditions. These plants are being used for the treatment of many human ailments including diabetes. Plants that are specifically employed for the treatment of diabetes are *Acacia nilotica*, *Acacia senegal*, *Aegle marmelos*, *Calotropis procera*, *Capparis deciduas*, *Cassia auriculata*, *Cassia sophera*, *Cayratia trifolia*, *Cyanopsis tetragonoloba*, *Dalbergia sisso*, *Gymnema sylvestre*, *Momordica charantia*, *Syzygium cumini*, *Withania somnifera*. This article aims to provide a comprehensive review on the some plants of Rajasthan having antidiabetic potential.

KEYWORDS: Diabetes mellitus, antidiabetic, ethnomedicine

INTRODUCTION

Diabetes mellitus is one of the most common chronic diseases in nearly all countries¹. This disease is characterized by hyperglycemia resulting from reduced (or absent) secretion of insulin, often coupled with reduced sensitivity to its action (insulin resistance)².

Diabetes mellitus fall into three broad categories: Type 1, Type 2, and Gestational diabetes. Type 1 diabetes mellitus is characterized by loss of insulin producing β -cells of the islets of Langerhans in the pancreas leading to insulin deficiency. This type of diabetes can affect children or adults and is traditionally termed "juvenile diabetes" as it represents a majority of the diabetes cases in children³. This type is less common and has lower degree of genetic predisposition. Type 2 diabetes mellitus is the most common type and characterized by insulin resistance which may be combined with relatively reduced insulin secretion. This type has higher degree of genetic predisposition and generally has a late onset (past middle age). It may occur due to abnormalities in glucoreceptors of β -cells, reduced sensitivity of peripheral tissues to insulin, reduction in number of insulin receptors and down regulation of these receptors³.

GLOBAL STATUS OF DIABETES MELLITUS

There are an estimated 285 million adults with diabetes in 2010; this number will continue to increase globally due to an aging population, growth of population size, urbanization and high prevalence of obesity and sedentary lifestyle. Diabetes leads to both premature death and complications such as blindness, amputations,

renal disease, and cardiovascular diseases¹. India is a vast, heterogeneous country with an approximate population of 1.1 billion people, a complex socio-political history, and immense diversity of culture, dialects and customs. World Health Organization (WHO) reported 32 million people with diabetes residing in India, while the International Diabetes Federation (IDF) has reported an estimated 40.9 million for the same country. In either case, India remains the country with the highest absolute number of people with diabetes worldwide⁴.

American Diabetes Association (ADA) Standards of Medical Care in Diabetes recommend a comprehensive disease management program that includes lifestyle interventions and use of oral antidiabetic agents (OADs) for the treatment of Type 2 Diabetes mellitus. Diabetic patients may experience the additive and/or adverse effects or tolerability issues associated with various OADs. For instance, sulfonylureas have been associated with weight gain and hypoglycemia, and patients on biguanides report negative gastrointestinal effects, while higher rates of weight gain, edema and fractures have been reported with thiazolidinediones⁵. Sometimes, patients may eventually require parenteral therapy with insulin or a glucagon-like peptide-1 (GLP-1) agonist⁶. As the presently available disease management strategies possess many patient incompatibilities, there is a need to discover more effective and safer antidiabetic agents by utilizing the rich heritage of medicinal plants.

MEDICINAL PLANTS OF RAJASTHAN WITH ANTIDIABETIC POTENTIAL

Rajasthan, the second largest state of the India exhibits variety in physiognomy, climate, soil and consequently the vegetation including a wide variety of medicinal plants⁷. The tribals of remote areas of Rajasthan are totally dependent on indigenous system of medicine for their healthcare as it is difficult for them to get modern medical facilities for their day-to-day health problems. The traditional healers of Rajasthan are having a commendable knowledge of the medicinal virtues of plants that grow around them⁸. Various ethnobotanical studies carried out in Rajasthan indicate that some plants (Table 1) are being used for their antidiabetic potential by most of the tribals; also prescribed by local vaid and bhopas (village priest).

Acacia nilotica (L.) Del. (Mimosaceae)

Local Names: Babul, Bambul

It is a tree that can grow up to the height of 10 m. The gum obtained from the stem is locally used in confectionery and sweets. Bhils take this gum orally to cure diabetes⁹.

Pharmacological Finding

1. A study was carried out to investigate the effect of an aqueous methanolic extract of pods on various biochemical parameters in alloxan-induced diabetic rabbits. Dose of 400 mg/kg body weight maximally reduced the blood glucose levels as compared to the diabetic group ($p < 0.001$). This dose also significantly ($p < 0.05$) lowered the plasma total cholesterol, triglyceride and low-density lipids (LDLs) in treated rabbits as compared to diabetic rabbits. This study revealed the antidiabetic potential of this plant¹⁰.

Acacia senegal (L.) Willd. (Mimosaceae)

Local Names: Kumat, Kumatio, Kumta

It is a small deciduous tree having smooth bark, pale-greenish grey in colour that peels off in flakes in older twigs¹¹. It is found on the hillsides and gravel throughout the state¹². The tree is extremely hardy and resistant to draught and is one of the main cash crops of the desert region¹³.

This tree yields the true gum arabic which is used as binding agent in the manufacture of cough pastilles in pharmaceutical industries and for its demulcent and emollient action¹⁴. The Bhopas in Udaipur region prescribe the gum as a part of food for diabetic patients⁹. However, the antidiabetic potential of this plant has not been evaluated scientifically.

Aegle marmelos (L.) Corr. (Rutaceae)

Local Names: Bel, Bel-patra

It is a small or medium sized tree that is sparsely scattered in mixed deciduous forests on the Aravalli

range in Rajasthan and also often planted all over India and frequently near the Hindu temples^{9, 15, 16}. The Garasia tribal take powder orally with water that is made up of roots, stem-bark and leaves to cure diabetes⁹.

Pharmacological Findings

1. A study was carried out to evaluate the antidiabetic as well as the antioxidant potential of methanolic extract of leaves. In this study, alloxan induced diabetic rats were fed with the extract (suspended in distilled water) daily at the dose of 100 mg/kg body weight. Oxidative stress produced by alloxan was found to be significantly lowered by the administration of this extract. Results indicated that methanolic extract of leaves effectively reduced the blood sugar level as well as oxidative stress in experimental animals¹⁷.

2. In another study, methanolic extract of the roots was fractionated into eight fractions using column chromatography. Antidiabetic activity of all the fractions was studied using the glucose uptake by isolated rat hemi-diaphragm *in vitro* model. The fractions coded as FB and FC enhanced the uptake of glucose by isolated rat hemi-diaphragm significantly ($p < 0.001$) and was found to be more effective than insulin.

Using the bioassay-guided fractionation, two compounds 1 and 2 (Figure 1) were isolated from fraction FC by column chromatography and identified as 6-methyl-4-chromanone and skimmianine respectively by NMR and mass spectral methods¹⁸.

3. Aqueous extract of seeds was administered orally at different doses (100, 250 and 500 mg/kg) to normal as well as sub (fasting blood glucose (FBG) normal; glucose tolerance abnormal) and mild (FBG 120–250 mg/dl) streptozotocin induced diabetic rats. The dose of 250 mg/kg was found to be the most effective dose and it decreased the blood glucose level (BGL) by 35.1% in normal healthy rats after 6 h of administration. The same dose also showed a marked reduction in BGL by 41.2% in sub and by 33.2% in mild diabetic rats in glucose tolerance test (GTT) after 2 h of administration. These results clearly indicated that aqueous seed extract possess antidiabetic effect in diabetic rats¹⁹.

Calotropis procera (Aiton) R. Br. (Asclepiadaceae)

Local Names: Madar, Aak, Aakro

It grows in the form of erect shrub or small tree and commonly found in wastelands and abundant in the desert throughout the year⁹. The tribals of Udaipur drink its root extract to cure diabetes and fever⁹.

Pharmacological Finding

Petroleum ether, methanolic and aqueous extracts of leaves and roots were investigated for its hypoglycemic effect in streptozotocin induced diabetic male wister albino rats. The different extracts were administered at

dose of 250 mg/kg, as single dose per day to the experimental animals for a period of 15 days. The blood glucose level serum and lipid profile was determined in the diabetic rats. This investigation established the pharmacological evidence for antidiabetic folklore claim²⁰.

***Capparis decidua* Linn. (Capparidaceae)**

Local Names: Ker, Kario

It is a much branched, xerophytic shrub that grows up to 15 feet height²¹. It is common throughout the state and generally found on the rocks, gravel and sandy plains. Sometimes, it occurs on sand-dunes forming the chief feature of the landscape¹². Unripe fruits are not only pickled locally, but also commercially. The mature fruits are eaten by the tribals. According to ethnopharmacological relevance, the dried fruits of *Capparis decidua* are used as an ingredient in antidiabetic compositions⁹.

Pharmacological Finding

In a study, alkaloid rich fraction (AR) was obtained from dried fruits and evaluated for antidiabetic potential in streptozotocin induced diabetic mice at a dose of 50 mg/kg body weight for 28 days. On completion of the treatment, it significantly inhibited the acute elevation of blood glucose level and also reduced total cholesterol (TC) and triglyceride (TG) content ($p < 0.05$). Activity of glucose-6-phosphatase (G6Pase) was attenuated by 44% and liver and muscle glycogen content showed significant improvement ($p < 0.05$). The expression of different target genes like G6Pase, phosphoenolpyruvate carboxykinase (PEPCK), aldose reductase and tumor necrosis factor- α (TNF- α) showed significant reduction whereas glucose transporter-4 (Glt-4), peroxisome proliferators activated receptor- γ (PPAR- γ) and glucokinase (GK) improved remarkably. This study clearly established the candidacy of AR fraction for further purification and characterization of the individual alkaloids, in order to understand their mechanism of action²².

***Cassia auriculata* L. (Caesalpinaceae)**

Local Names: Anwal, Tarwan, Tarawar

It is a tall, much branched shrub that grows up to 2 m height having reddish-brown smooth bark. It is found in gravelly and sandy soil throughout the state. Its bark provides one of the most important tanning materials¹². Bhils take orally the fresh flowers with sugar to cure diabetes^{9, 23}. Its seeds and leaves are also used in diabetes⁹.

Pharmacological Findings

1. Leaf extract of this plant was studied for hypoglycemic activity in alloxan induced fasted diabetic rats and effect of this extract was also studied for glucose

utilization by isolated rat hemi diaphragm method. For this purpose, ethanolic and water extract at a dose of 200 mg/kg per day was administered to diabetic rats for 10 consecutive days. A significant reduction in serum glucose level was observed from 3rd day to till the end of the experiment. During another study, the albino rats were killed by decapitation and diaphragms were exposed to Group I; tyrode solution with glucose (2000 mg/l) only, served as control, Group II; tyrode solution with glucose (2000 mg/l) + Insulin (0.25 IU/ml), Group III; tyrode solution with glucose (2000 mg/l) + extract of *C. auriculata* (25 mg/ml) and group IV; tyrode solution with glucose (2000 mg/l) + Insulin (0.25 IU/ml) + extract of *C. auriculata* (25 mg/ml). Glucose uptake and glycogen deposition studies suggested that leaf extract probably has no direct insulin like effect which can enhance the peripheral utilization of glucose²⁴.

2. Aqueous extract of leaves was evaluated for antihyperglycemic and hypolipidemic activity in streptozotocin induced mild diabetic (MD) and severe diabetic (SD) rats. The extract was orally administered to MD and SD rats at 100, 200 and 400 mg/kg doses for 1 day to determine antihyperglycemic activity. The 400 mg/kg dose was administered daily for 3 weeks to assess glycemic control and hypolipidemic effect. A dose dependant fall in fasting blood glucose was observed after 5 h of extract administration in diabetic rats. After 3 weeks treatment, extract produced significant reduction in fasting blood glucose and glycosylated haemoglobin in both MD and SD rats. Serum lipid levels were reversed towards normal in extract fed MD and SD rats. It demonstrated that this plant possesses potent antihyperglycemic and hypolipidemic activity in both MD and SD rats²⁵.

***Cassia sophera* L. (Caesalpinaceae)**

Local Name: Kasunda

It is a shrub that is found chiefly in the east of Aravalli and rarely found in wastelands. Bhils of eastern Rajasthan mix its bark and seed powder in honey and give to the diabetic patients to eat^{9, 26}.

Pharmacological Finding

Aqueous and methanol extracts of seeds has been reported to possess significant hypoglycemic activity in alloxan induced diabetic rabbits²⁷.

***Cayratia trifolia* (L.) Domin (Vitaceae)**

Local Names: Char, Char-bel, Kihata nimba

It is a climbing or spreading herb or undershrub with 3-foliolate leaves^{9, 28, 29}. It is easily found in Sitamata wildlife sanctuary of Chittorgarh and Udaipur district located in south-west region of the state²⁸. Its stem, leaves and root contain hydrocyanic acid and leaves also contain delphinidin, cyanidin and yield yellow waxy oil and

sterols¹³. Bhils of Udaipur region take orally the extract of roots to cure diabetes⁹. The extract of tuber of this plant along with the infusion of its seeds is given orally to diabetic patients to check the sugar level of blood²⁵. However, the antidiabetic potential of *Cayratia trifolia* has not been evaluated scientifically.

***Cyamopsis tetragonoloba* (L.) Taub. (Fabaceae)**

Local Names: Guar, Ganwar, Gwar

It is an annual herb that is cultivated as a rainy season crop for edible fruits, seeds and fodder throughout the state^{9,13}. The young pods are consumed as vegetable and whole plant is used as a green manure¹². Tribals of the desert soak the seeds in water in the evening and drink that water in the next morning to cure diabetes⁹.

Pharmacological Finding

Aqueous extract of beans was investigated for fasting blood glucose levels in glucose loaded normal and alloxan induced diabetic rats. Aqueous extract of beans at 250 mg/kg body weight significantly lowered blood glucose levels in diabetic rats within 3 hr of administration. Continued administration of the extract at the same dose daily for 10 days produced statistically significant reduction in the blood glucose levels indicating the hypoglycemic potential of beans of this plant³⁰.

***Dalbergia sisso* Roxb. (Fabaceae)**

Local Name: Sisham

It is a deciduous tree that is common in wastelands, along the roads and boundaries of gardens⁹. It yields a valuable timber that has high industrial and commercial value^{9,12}. Tribals of Dungarpur district take orally the paste of leaves to cure diabetes⁹.

Pharmacological Finding

In a study, the ethanolic extract of leaves was administered orally at the doses of 250 and 500 mg/kg to normal rats. Dose of 500 mg/kg was found to be more effective; it decreased blood glucose level (BGL) by 38.2 % in normal healthy rats after 1 day of administration. After daily treatment with both the doses of extract for 21 days to alloxan induced diabetic (FBG 300-350 mg/dL) rats, the BGL reduced to 125 mg/dL by 250 mg/kg and 104 mg/dL by 500 mg/kg. This study indicated the hypoglycemic and antihyperglycemic potential of the extract³¹.

***Gymnema sylvestre* (Retz.) R. Br. (Asclepidaceae)**

Local Names: Jangli-urad, Gurmar

It is a much branched, twining woody climber running over the tops of small trees³². It is frequently found in the deciduous forests in the east of Aravalli⁹. Its leaf extract contain gymnemic acid which is said to inhibit hyperglycemia. It abolishes the taste of sugar and is believed to neutralize the excessive sugar present in the

body of diabetic patient³². It is used as a remedy for diabetes throughout the state⁹.

Pharmacological Findings

1. In a study, aqueous extract of leaves was administered orally at the dose of 400, 600 and 800 mg/kg for 30 days. Fasting blood glucose, cholesterol, HDL cholesterol and serum triglyceride levels were estimated in both normal and alloxan induced diabetic rats. All these levels were found to be significantly reduced in treated rats. It revealed that this plant possess significant antidiabetic and hypolipidemic activity and seems promising for the development of a phytomedicine for treatment of diabetes mellitus³³.

2. In another study, an active compound-dihydroxy gymnemic triacetate (Figure 2) has been isolated from its acetone extract. This compound at a dose of 20 mg/kg body weight was administered orally for 45 days to streptozotocin diabetic rats for the assessment of various biochemical parameters in normal and diabetic rats. This compound produced significant effects on all biochemical parameters in comparison to diabetic control group. Results indicated the hypoglycemic and hypolipidemic activity of dihydroxy gymnemic triacetate on long-term treatment and it could be used as a drug for treating diabetes³⁴.

***Momordica charantia* L. (Cucurbitaceae)**

Local Names: Karela, karelo

It is a monoecious climber and cultivated for its fruits cooked as vegetable. Local vaid prescribe its vegetable to the diabetic patients⁹. It is a medicinal plant and used in Ayurveda for treating various diseases including diabetes mellitus³⁵.

Pharmacological Findings

1. Aqueous extract of fresh unripe whole fruits at a dose of 20 mg/kg body weight was found to reduce fasting blood glucose by 48%, comparable to the standard drug-glibenclamide. This extract was tested for nephrotoxicity, hepatotoxicity and biochemical parameters such as SGOT, SGPT and lipid profile. The extract did not show any signs of nephrotoxicity and hepatotoxicity as judged by histological and biochemical parameters. Thus the aqueous extract was proved as a safe alternative for reducing blood glucose level³⁰.

2. In vitro and in vivo experiments were carried out to study the role of juice of fruits on the diabetic status. Control and streptozotocin induced diabetic mice were fed orally daily once with either 10, 50 or 100% juice (0.2 ml juice equivalent to 10 ml/kg body weight) for a period of 5 days. Activity of the juice was tested on streptozotocin treated RIN cells (rat insulinoma cell line) and isolated islets in vitro. Feeding with juice caused reduction in streptozotocin-induced hyperglycemia in

mice. Further it also reduced the streptozotocin-induced apoptosis in RIN cells indicating the mode of protection of RIN cells, islets and pancreatic β -cells¹⁵.

3. Acetone extract of whole fruit was studied at the doses of 25, 50 and 75 mg/100 g body weight in alloxan induced diabetic rats. Histological observations showed the different phases of recovery of β -cells of the islets of Langerhans of pancreas, which were less in number in the untreated diabetic rats proving the antidiabetic potential of this plant¹⁷.

***Syzygium cumini* (L.) (Myrtaceae)**

Local Names: Jamun, Jambus

It is an evergreen tree having grayish brown branches²¹. Local vaidas prescribe the mature fruits to the diabetic patients⁹. Leaves, seeds, wood and stem bark are also used in treatment of diabetes¹². An aqueous extract of seeds found to cause the marked, prolonged decrease in blood sugar upon hypodermic injection into dogs¹⁶.

Pharmacological Finding

Oral administration of petroleum ether, chloroform, acetone, methanolic, and water extracts at a dose of 100 mg/kg, p.o. for 21 days caused a decrease in fasting blood sugar (FBS) in diabetic rats. Among all the extracts, methanolic extract was found to lower the FBS significantly in diabetic rats. This extract was subjected to column chromatography that led to the isolation of an active principle, Cuminoside (a phenolic glycoside). It was studied for its hypoglycemic and antioxidant potential at the dose of 50 mg/kg, p.o. and a significant decrease in FBS level, lipid peroxidation level, and improvement in the levels of antioxidant enzymes was observed in diabetic rats. This study provided a scientific rationale for the use of cuminoside as an antidiabetic agent²³.

***Withania somnifera* (L.) (Solanaceae)**

Local Names: Ashgandh, Chirpotan, Rashbhari

It is a much branched, erect, perennial undershrub that grows up to the height of 1.5 m²⁰. It is common in moist and shady habitats^{9,12}. It is an Ayurvedic medicinal plant that is popular home remedy for several human ailments. It is mentioned in Vedas as an herbal tonic and health food. Infusion of leaves is prescribed by local vaidas as a medicine to cure diabetes⁹. Its roots also exhibit hypoglycemic activity²².

Pharmacological Findings

1. Hypoglycaemic and hypolipidaemic effects of extracts of root and leaves were investigated at doses of 100 and 200 mg/kg body weight in alloxan induced diabetic rats. After 8 weeks of treatment with the extract, a significant decrease was observed in the levels of urine sugar, blood glucose, glycosylated haemoglobin, liver glucose-6-phosphatase, aspartate transaminase, alanine

transaminase, acid phosphatase, alkaline phosphatase, serum lipids except high density lipoprotein-bound cholesterol. It indicated the hypoglycemic and hypolipidaemic activity of root and leaf extract of this plant¹⁹.

2. Hypoglycemic, diuretic and hypocholesterolemic effects of roots were assessed on human subjects. 6 mild NIDDM subjects and 6 mild hypercholesterolemic subjects were treated with the powder of roots for 30 days. Various parameters were studied in the blood and urine samples of the subjects along with dietary pattern before and at the end of treatment period. Decrease in blood glucose was comparable to that of standard drug. Significant increase in urine sodium, urine volume, significant decrease in serum cholesterol, triglycerides, LDL and VLDL were observed and indicated the hypoglycemic, diuretic and hypocholesterolemic activities of root⁴⁰.

CONCLUSION

Research studies have a role in highlighting inequalities and improving comprehension of the complex mix of predisposing and modifiable factors associated with a particular disease. The research in alternative system of medicine indicates that further exploration of natural resources is required to establish the pharmacological activities and to find the active constituents for the development of drugs. These studies provide the right blend of evidence to stimulate the modification of current medical practices. Geno- and phenotypically, the Asian Indian population are at elevated risk for developing diabetes in comparison to other population groups. Therefore, the well planned, dedicated, and diligent care should be a priority for diabetes patients. Plant origin based drugs have been utilizing from many years in patients with insulin dependent and non-insulin dependent diabetes. Scientific evaluations of the plant species have proved their efficacy for serving as the hypoglycemic agent. Conclusively, the integration of research in alternative system of medicine and clinical practice is must in seeking sustainable, practical, scientific, and socially acceptable models for diabetes care.

REFERENCES

1. Zhang P, Zhang X, Brown J, Vistisen D, Sicree R, Shaw J et al. Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes Res Clin Pract.* 2010; 87:293-301.
2. Rang HP, Dale MM, Ritter JM, Moore PK. *Pharmacology.* 5th ed. New Delhi: Elsevier; 2003
3. Tripathi KD. *Essentials of Medical Pharmacology.* 4th ed. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd.; 2001
4. Ali MK, Narayan KMV, Mohan V. Innovative research for equitable diabetes care in India. *Diabetes Res Clin Pract.* 2009; 86: 155-167.

5. Pollack MF, Purayidathil FW, Bolge SC, Williams SA. Patient-reported tolerability issues with oral antidiabetic agents: Associations with adherence, treatment satisfaction and health-related quality of life. *Diabetes Res Clin Pract.* 2010; 87: 204-210.
6. Rodbard HW, Green AJ, Fox KM, Grandy S. Impact of type 2 diabetes mellitus on prescription medication burden and out-of-pocket healthcare expenses. *Diabetes Res Clin Pract.* 2010; 87: 360-365.
7. Singh VK, Govil JN, Singh G. Recent Progress in Medicinal Plants: Ethnomedicine and Pharmacognosy. Vol 1. Houston, Texas: SCI TECH Publishing LLC; 2002
8. Jain A, Katewa SS, Galav P, Nag A. Some therapeutic uses of biodiversity among the tribals of Rajasthan. *Indian Journal of Traditional Knowledge.* 2008; 7(2): 256-262.
9. Singh V, Pandey RP. Ethnobotany of Rajasthan, India. 1st ed. Jodhpur: Scientific Publishers (India); 1998
10. Ahmad M, Zaman F, Sharif T, Zabta M. Antidiabetic and Hypolipidemic Effects of Aqueous Methanolic Extract of *Acacia Nilotica* Pods in Alloxan-Induced Diabetic Rabbits. *Scandinavian Journal of Laboratory Animal Science.* 2008; 35(1): 29-34.
11. Sharma S, Tiagi B. Flora of North-East Rajasthan. New Delhi: Kalyani Publishers; 1979
12. Bhandari MM. Flora of the Indian Desert. 1st ed. Jodhpur: MPS Reprints; 1978 (Revised ed. 1990)
13. The Wealth of India. Vol 3. New Delhi: Publications & Information Directorate, CSIR; 1992
14. Kashyapa K, Chand R, editors. The Useful Plants of India. 1st ed. New Delhi: National Institute of Science Communication; 1986 (4th Reprint 2000)
15. Kartikar KR, Basu BD. Indian Medicinal Plants. Vol. I, 2nd ed. Dehradun, India: International Book Distributors; Reprint 1987
16. Chopra RN, Nayar SL, Chopra IC. Glossary of Indian Medicinal Plants. 1st ed. New Delhi: National Institute of Science Communication (CSIR); 1956 (5th Reprint 1999)
17. Sabu MC, Kuttan R. Antidiabetic activity of *Aegle marmelos* and its relationship with its antioxidant properties. *Indian J Physiol Pharmacol.* 2004; 48(1): 81-88.
18. Ravi S, Sadashiva CT, Tamizmani T, Balasubramanian T, Rupeshkumar M, Balachandran I. In vitro glucose uptake by isolated rat hemi-diaphragm study of *Aegle marmelos* Correa root. *Bangladesh J Pharmacol* 2009; 4: 65-68.
19. Kesari AN, Gupta RK, Singh SK, Diwaker S, Watal G. Hypoglycemic and antihyperglycemic activity of *Aegle marmelos* seed extract in normal and diabetic rats. *J Ethnopharmacol* 2006; 107: 374-379.
20. Bhaskar VH, Singh S. Evaluation of antihyperglycemic activity of extracts of *Calotropis procera* (ait.) R.br on streptozotocin induced diabetic rats. *Global Journal of Pharmacology* 2009; 3(2): 95-98.
21. Bentley R, Trimen H. Medicinal Plants, Vol 3. 2nd Indian Reprint. Dehradun: Allied Book Centre; 1992
22. Sharma B, Salunke R, Balomajumder C, Daniel S, Roy P. Anti-diabetic potential of alkaloid rich fraction from *Capparis decidua* on diabetic mice. *J Ethnopharmacol* 2010; 127: 457-462.
23. Kartikar KR, Basu BD. Indian Medicinal Plants. Vol. II, 2nd ed. Dehradun, India: International Book Distributors; Reprint 1987
24. Sabu MC, Subburaju T. Effect of *Cassia auriculata* Linn. on serum glucose level, glucose utilization by isolated rat hemidiaphragm. *J Ethnopharmacol* 2002; 80: 203-206.
25. Gupta S, Sharma SB, Bansal SK, Prabhu KM. Antihyperglycemic and hypolipidemic activity of aqueous extract of *Cassia auriculata* L. leaves in experimental diabetes. *J Ethnopharmacol* 2009; 123: 499-503.
26. Nadkarni KM. Indian Plants and Drugs with their Medical Properties and uses. Reprint Indian ed. New Delhi: Asiatic Publishing House; 2001
27. Khan AM, Khan AH, Akhtar MS, Ahmad B, Sher A, Ahmed W. Pharmacological screening of *Cassia sophera* for hypoglycemic activity in normal and Diabetic Rabbits. *Proceedings-Shaikh Zayed Postgraduate Medical Institute* 2002; 16(1):1-4.
28. Jain A, Katewa SS, Galav PK, Sharma P. Medicinal plant diversity of Sitamata wildlife sanctuary, Rajasthan, India. *J Ethnopharmacol* 2005; 102: 143-157.
29. Swarnkar S, Katewa SS. Ethnobotanical observation on tuberous plants from tribal area of Rajasthan (India). *Ethnobotanical leaflets* 2008; 12: 647-666.
30. Mukhtar HM, Ansari SH, Ali M, Bhat ZA, Naved T. Effect of aqueous extract of *Cyamopsis tetragonoloba* Linn. beans on blood glucose level in normal and alloxan-induced diabetic rats. *Indian J Exp Biol.* 2004; 42(12): 1212-1215.
31. Singh P, Singh D, Prajapati K, Jain SK. Antidiabetic activity of ethanolic extract of *Dalbergia sissoo* L. leaves in alloxan induced diabetic rats. *International Journal of Current Pharmaceutical Research* 2010; 2(2): 24-27.
32. Khan IA, Khanum A. Medicinal and Aromatic Plants of India. 1st ed. Hyderabad, Andhra Pradesh: Ukaaz Publications; 2005. p. 90, 197.
33. Mall GK, Mishra PK, Prakash V. Antidiabetic and hypolipidemic activity of *Gymnema sylvestre* in alloxan induced diabetic rats. *Global Journal of Biotechnology & Biochemistry* 2009; 4(1): 37-42.
34. Daisy P, Eliza J, Farook KAMM. A novel dihydroxy gymnemic triacetate isolated from *Gymnema sylvestre* possessing normoglycemic and hypolipidemic activity on STZ-induced diabetic rats. *J Ethnopharmacol* 2009; 126: 339-344.
35. Sitasawad SL, Shewade Y, Bhonde R. Role of bittergourd fruit juice in stz-induced diabetic state in vivo and in vitro. *J Ethnopharmacol* 2000; 73: 71-79.
36. Virji J, Sivakami S, Shahani S, Suthar AC, Banavalikar MM, Biyani MK. Antihyperglycemic effects of three extracts from *Momordica charantia*. *J Ethnopharmacol* 2003; 88: 107-111.
37. Singh N, Gupta M. Regeneration of β cells in islets of Langerhans of pancreas of alloxan induced diabetic rats by acetone extract of *Momordica charantia* (Linn.) (bitter gourd) fruits. *Indian J Exp Biol.* 2007; 45: 1055-1062.
38. Farswan M, Mazumder PM, Partha V, Upaganlawar A. Modulatory Effect of *Syzygium cumini* Seeds and its Isolated Compound on biochemical parameters in Diabetic Rats. *Pharmacogn Mag.* 2009; 5(18): 127-133.
39. Udayakumar R, Kasihurirengan S, Mariashibu TS, Rajesh M, Anbazhagan VR, Kim SC et al. Hypoglycaemic and Hypolipidaemic Effects of *Withania somnifera* Root and Leaf Extracts on Alloxan-Induced Diabetic Rats. *Int J Mol Sci* 2009; 10: 2367-2382.
40. Andallu B, Radhika B. Hypoglycemic, diuretic and hypocholesterolemic effect of winter cherry (*Withania somnifera*, Dunal) root. *Indian J Exp Biol* 2000; 38(6):607-9.

TABLE 1: IT SHOWS MEDICINAL PLANTS OF RAJASTHAN WITH ANTIDIABETIC POTENTIAL

S. No.	Name of plant, Family	Part used	Extract/Active constituent
1.	<i>Aracia nilotica</i> L. Del., Mimosaceae	Gum, stem bark, pods	Aqueous-methanol extract
2.	<i>Acacia senegal</i> L. Walld., Mimosaceae	Gum	--
3.	<i>Aegle marmelos</i> L. Corr., Rutaceae	Fruits, stem bark, root, leaves	Aqueous & methanol extracts/ 6-methyl-4-chromanone, skimmianine
4.	<i>Crotalaria procera</i> (Aiton) R. Br., Asclepiadaceae	Root, leaves	Petroleum ether, aqueous & methanol extracts
5.	<i>Capparis decidua</i> Lam., Cappariaceae	Fruits, stem bark, flower	Alcoholic extract/ Alkaloid rich fraction
6.	<i>Cassia aspiculata</i> L., Caesalpinaceae	Leaves, seeds, flower	Aqueous extract
7.	<i>Cassia sophera</i> L., Caesalpinaceae	Stem bark, seeds	Aqueous & methanol extracts
8.	<i>Cyrtaria trifolia</i> L. Donin, Vitaceae	Root, seeds	Infusion
9.	<i>Cyamopsis tetragonoloba</i> L. Taub., Fabaceae	Gum, pods	--
10.	<i>Dalbergia eisso</i> Roxb. Fabaceae	Leaves	Ethanol extract
11.	<i>Gymnema sylvatica</i> (Ratz.) R. Br., Asclepiadaceae	Leaves	Aqueous extract/ Dihydroxy gymnemaic triacetate
12.	<i>Misandica charantia</i> L., Cucurbitaceae	Fruits	Acetone, alcoholic & aqueous extracts
13.	<i>Syzygium cumini</i> L., Myrtaceae	Fruits, stem bark, seeds, leaves	Methanol extract/ Cuminoic acid (phenolic glycoside)
14.	<i>Withania somnifera</i> L., Solanaceae	Leaves, root	Alcoholic extract




Figure 1: It shows the structures of compound 1 (6-methyl-4-chromanone) and compound 2 (skimmianine)




Figure 2: It shows the structure of dihydroxy gymnemaic triacetate isolated from acetone extract of *Gymnema sylvatica*