



## EVALUATION OF ANTI-INFLAMMATORY ACTIVITY OF *NARAVELIA ZEYLANICA* LINN. LEAVES EXTRACT

Moses Samuel Rajan\*, Yedle Randhir, Shenoy Ashok, Sabraya A.R., Rahul G Raut  
Department of Pharmacology, Srinivas college of Pharmacy, Mangalore, Karnataka, India

Article Received on: 10/04/12 Revised on: 21/05/12 Approved for publication: 09/06/12

\*Email: randhir\_05@rediffmail.com

### ABSTRACT

*Naravelia Zeylanica* Linn. commonly known as Vatnisini is traditionally used for several medicinal purposes in India. The present study assessed anti-inflammatory activity of its leaf. Dried and crushed leaves of *Naravelia Zeylanica* were extracted with ethanol. The ethanol extract (EENZ) at the doses of 250 mg/kg and 500 mg/kg body weight was subjected for evaluation of anti-inflammatory activity in experimental animal models. Acute anti-inflammatory activity was evaluated by Carrageenan, Histamine and Serotonin-induced paw oedema in Wistar albino rats. Diclofenac sodium was employed as reference drug for anti-inflammatory study. In the present study, the ethanol extract of the leaves of *Naravelia Zeylanica* demonstrated significant acute anti-inflammatory activity in the tested models.

**Key words:** *Naravelia Zeylanica* Linn., Anti-inflammatory, Pain

### INTRODUCTION

Inflammation is a normal protective response to tissue injury caused by physical trauma, noxious chemical or microbial agents. It is the body response to inactivate or destroy the invading organisms, to remove irritant and set the stage for tissue repair. It is triggered by the release of chemical mediators from injured tissue and migrating cells<sup>1</sup>. Drugs from plant origin are used in India for treatment of many diseases in traditional system of medicine. *Naravelia zeylanica* DC (Ranunculaceae) is a climbing shrub with tuberous roots; wiry stem strong tendrils, leaves 3-foliate, opposite, terminal leaflet modified into a 3 branched tendril, leaflets ovate-lanceolate, serrate or crenate, prominently nerved; flowers yellow, fragrant, in axillary and terminal panicles, sepals downy, petals linear-clavate, elongate; fruits aggregate of achenes, ending in twisted feathery tails. The plant is available rich all around south India<sup>2</sup>. Plant is reported to contain mainly alkaloids, flavonoids, saponins and tannins. Methanolic extract of leaves of *Naravelia zeylanica* berberine is present, sterols are also present in the ethanolic extract of leaves and stem<sup>3</sup>. The whole plant is traditionally used in vitiated vata, pitta, inflammation, skin diseases, arthritis, headache, colic, wounds and ulcers. Leaf paste is consumed to treat Chest pain. The vines when crushed give a pungent odour which is inhaled to cure cold, all type of headaches including migraine<sup>3</sup>. Root and stem paste is applied externally for psoriasis, itches and skin allergies. The traditional medicine practitioners using the leaf and stem juices for treating intestinal worms, psoriasis & dermatitis<sup>4</sup>. The literature survey reveals no reports on the anti-inflammatory activity of the leaves extracts of *Naravelia Zeylanica*. This prompted us to investigate the anti-inflammatory activity of *Naravelia Zeylanica* leaves extract.

### MATERIALS AND METHODS

#### Plant material

The leaves of *Naravelia Zeylanica* were collected from Udupi, Karnataka, during July. It was authenticated by Dr. Gopalakrishna Bhat, Department of Botany, Poorna Prajna College, Udupi, Karnataka, India. A voucher specimen has been maintained in our lab for future reference.

#### Preparation of Extract

The leaves of *Naravelia zeylanica* were collected from Udupi, Karnataka. Fresh leaves of *N. zeylanica* were shade dried, powdered mechanically & stored in airtight containers until the extraction. The powder extracted with 95% ethanol in Soxhlet extractor for 30 hour. Extract was concentrated under reduced pressure using rotary flash evaporator to a syrupy consistency. Then it was dried in the desiccator (68% yield).<sup>5</sup>

**Animal Used:** Wistar strain rats of either sex (150-180 gm) of either sex were procured from Indian Institute of Sciences. They are maintained under standard conditions (temperature  $22 \pm 2^\circ\text{C}$ , relative humidity  $50 \pm 5\%$  and 12 hour light/dark cycle). The animals were housed in sanitized polypropylene cages containing sterile paddy husk as bedding. They had free access to standard pellet diet and water *ad libitum*. The Institutional Animal Ethics Committee approved the experimental protocol (Ref no. 367001/C/CPCACA). All the animals received humane care according to the criteria outlined in the "Guide for the Care and Use of Laboratory Animals" prepared by the "National Academy of Sciences" and published by the "National Institute of Health".

#### Evaluation of anti-inflammatory activity

##### Carrageenan induced rat paw oedema

The rats were divided into four groups containing six rats in each group. 0.1 ml of 1.0% carrageenan in normal saline (0.9% w/v NaCl) was injected to the sub plantar region of right hind paw. The EENZ was administered to the rats 1 h before carrageenan injection. Different groups were treated as follows:

Group I: Carrageenan (0.1 ml of 1.0% carrageenan/rat to the sub plantar region).

Group II: Carrageenan + Diclofenac sod. (10 mg/kg b. w., p. o.)

Group II and IV: Carrageenan + EENZ (250 mg/kg and 500 mg/kg b. w., p. o. respectively).

The paw volume was measured initially and at 0.5, 1, 3 and 5 h after carrageenan injection, using Plethysmograph, inflammation was calculated for comparison<sup>6,7</sup>.

**Histamine induced rat paw oedema**

The rats were divided into four groups containing six rats in each group. 0.1 ml of 1.0% histamine sulphate in normal saline (0.9% w/v NaCl) was injected to the sub-plantar region of right hind paw. The EENZ was administered to the rats 1 h before histamine injection. Different groups were treated as follows:

Group I: Histamine (0.1 ml of 1.0% histamine/rat to the sub plantar region).

Group II: Histamine + Diclofenac sod. (10 mg/kg b. w., p. o.)

Group III and IV: Histamine + EENZ (250 mg/kg and 500 mg/kg b.w., p.o. respectively).

The paw volume was measured initially and at 0.5, 1, 3 and 5 h after histamine injection, using Plethysmograph, inflammation was calculated for comparison<sup>6,7</sup>.

**Serotonin induced rat paw oedema**

The rats were divided into four groups containing six rats in each group. 0.1 ml of 1.0% serotonin in normal saline (0.9% w/v NaCl) was injected to the sub plantar region of right hind paw. The EENZ was administered to the rats 1 h before serotonin injection. Different groups were treated as follows:

Group I: Serotonin (0.1 ml of 1.0% serotonin/rat to the sub plantar region).

Group II: Serotonin + Diclofenac sod. (10 mg/kg b. w., p. o.)

Group III and IV: Serotonin + EENZ (250 mg/kg and 500 mg/kg b.w., p.o. respectively).

The paw volume was measured initially and at 0.5, 1, 3 and 5 h after serotonin injection, using Plethysmograph, inflammation was calculated for comparison<sup>6,7</sup>.

**Cotton pellet induced granuloma**

This is sub-acute model for inflammatory study. This method was adopted from D'Arcy (1960), which was carried out by using sterilized cotton pellet implantation method in rats. Under light ether anaesthesia, using blunted forceps subcutaneous tunnel was made and sterilized cotton pellets (10 ± 1 mg) were implanted in the axilla and groin region of the rat. Different groups were treated as follows:

Group-I: Vehicle control received 1% Tween-20 (dose: 10 ml/kg).

Group-II: Animals treated with Diclofenac sod (dose: 10 mg/kg).

Group-III and IV: Animals treated with + EENZ (250 mg/kg and 500 mg/kg b.w., p.o. respectively). After recovering from anaesthesia, animals were treated orally with vehicle control, diclofenac sod. and various doses of herbal formulation for consecutive 7 days, once per day. They were sacrificed on day 8th by cervical dislocation and the pellets were removed, freed from extraneous tissue and dried at 60°C for 24 hrs. The percentage inhibition of the dry weight of the granuloma were calculated and compared.<sup>8</sup>

**Statistical Analysis**

All data were expressed as the mean ± SEM. The results were analyzed for statistical significance (P<0.05) by One-way (ANOVA) followed by Dunnett's test using computerized Graph Pad InStat version 3.05, Graph pad software, U.S.A.

**RESULT AND DISCUSSION**

The present study evaluated the anti-inflammatory activity of ethanolic extract from *N. Zeylanica* leaves in experimental rodent models. The results are presented in Tables 1-4.

In Indian system of medicine, certain herbs are claimed to provide relief of pain and inflammation. The claimed therapeutic reputation has to be verified in a scientific manner. In the present study one such drug *N. Zeylanica* was

taken for the study. The leaves extracts of *N. Zeylanica* possess significant anti-inflammatory effect in the anti-inflammatory model of inflammation in rats.

The carrageenan, Histamine and serotonin-induced hind paw oedema a model in rats are known to be the acute inflammatory model sensitive to cyclooxygenase (COX) inhibitors and has been used to evaluate the effect of nonsteroidal anti-inflammatory agents (NSAID), which primarily inhibit the cyclooxygenase involved in prostaglandin (PG) synthesis. In case of the time course of oedema development in these paw oedema models in rats are generally two phases are found. The first phase, which occurs between 0 to 2.5 hour of injection of the phlogistic agent, has been attributed to the release of histamine or serotonin. The oedema volume reaches to its maximum approximately 3 hour post treatment and then begin to decline. The second phase of inflammatory reaction which is measured at 3hour is caused by the release of bradykinin, protease, prostaglandin and lysosome. Therefore, it can be inferred that the inhibitory effect of the extract on the carrageenan, Histamine and Serotonin induced inflammation could be due to the inhibition of enzyme cyclooxygenase leading to inhibition of prostaglandin synthesis. Thus, the results of the present study demonstrates that the ethanolic extract obtained from leaves of *Naravelia Zeylanica* exhibited acute anti-inflammatory activity in the tested models which was found to be the most effective at higher concentrations employed.

The cotton pellet granuloma method has been widely employed to assess the transudative, exudative and proliferative components of chronic inflammation. The fluid absorbed by the pellet greatly influences the wet weight of the granuloma. The results indicate that extract of *N. Zeylanica* has anti-transudative and anti-proliferative activity. However, a more extensive study is necessary to determine the exact mechanism of action of the extract and its active compound.

**CONCLUSION**

It is concluded that Ethanolic extract of leaves of *Naravelia Zeylanica* possess significant anti-inflammatory activity against all experimental models in rats. This may be due to the presence of reported active Phyto-constituents & their influence on the prostaglandins pathway. Further research, to isolate anti-inflammatory principle & exact mechanism involved, is needed.

**ACKNOWLEDEMENT**

The authors are grateful to management of Srinivas college of Pharmacy, Mangalore for providing necessary facilities to carry out the experiments and A.Shama Rao Foundation, Mangalore for providing financial assistance.

**REFERENCES**

1. Bhitre, MJ, Fulmali S, Kataria M, Anwikar S, Kadri H. Antiinflammatory activity of the fruits of *piper longum* Linn. Asian J Chemistry 2008; 20(6): 4357-60.
2. Raja H, Naika B, Krishna V. Micropropagation, Isolation and characterization of Berberine from leaves of *Naravelia zeylanica* DC. Res J of Medicinal plant 2008; 2(1):1-9.
3. Raja HN, Krishna V, Harish BG, Khadeer A, Mahadevan KM. Antimicrobial Activity of Bioactive Constituents Isolated from Leaves of *Naravelia zeylanica*. Intl J of Biomed Pharma Sci 2002; 1:153-59.
4. Udayan PS, Satheesh G, Tushar KV, Indira B. Medicinal plants used by Malayali tribes of Sevarayan hills Yercad, Salem, Tamilnadu, India. Zoos Print J 2006; 21(4): 222-24.
5. Ashoka SM, Shastry CS, Sridevi, Gopkumar p. Antiulcer activity of *Naravelia zeylanica* leaves extract. J of Pharmacy Research 2009;2(7):1218-20.
6. Biswas M, Biswas, Ghosh AK, Halder PK. A pentacyclic triterpenoid

- possessing anti-inflammatory activity from the fruits of *Dregea volubilis*. Phcog Mag 2009; 5: 64-68.
7. Majumder AM, Naik DG, Dandgee CN, Puntambekar HM. Antiinflammatory Activity of *Curcuma amada* Roxb. In albino rats. Indian J Pharmacol 2000; 32: 375-377.
  8. Arcy D, Haward PF, Muggleton RW, Townsend SB. The Anti-inflammatory action of Griseofulvin in experimental animals. J of Pharmacy and Pharmacology 1960; 12: 659-665.
  9. Alcatraz MJ, Jimenez MJ. Flavonoids as anti-inflammatory agents. Fitoterapia 1998; 59: 25-38.
  10. Della LR, Tubaro PD, Zilli C, Del NP. The role of flavonoids in the anti-inflammatory activity of *Chamomilla recutita*. Clin Biol Res 1986; 213: 481-488.

**Table 1. Anti-inflammatory effect of EENZ on carrageenan induced rat paw oedema**

| Sr.no | Treatment       | Dose mg/kg p.o. | 0.5h       | 1h          | 3h           | 5h           | % Inhibition |
|-------|-----------------|-----------------|------------|-------------|--------------|--------------|--------------|
| 1     | Control         |                 | 0.21±0.020 | 0.41±0.018  | 0.43±0.012   | 0.47±0.015   | -            |
| 2     | Diclofenac sod. | 10              | 0.17±0.012 | 0.19±0.014* | 0.11±0.020** | 0.07±0.015** | 65.81        |
| 3     | EENZ            | 250             | 0.19±0.017 | 0.31±0.009  | 0.27±0.014*  | 0.18±0.015*  | 37.23        |
| 4     | EENZ            | 500             | 0.17±0.008 | 0.22±0.010* | 0.19±0.012*  | 0.11±0.013** | 56.26        |

Values are mean ± SEM (n = 6). \* $p < 0.05$ , \*\* $p < 0.001$  when compared to control

**Table 2. Anti-inflammatory effect of EENZ on Histamine induced rat paw edema**

| Sr no | Treatment       | Dose mg/kg p.o. | 0.5h        | 1h          | 3h          | 5h           | % inhibition |
|-------|-----------------|-----------------|-------------|-------------|-------------|--------------|--------------|
| 1     | Control         |                 | 0.35±0.020  | 0.31±0.014  | 0.27±0.02   | 0.33±0.020   | -            |
| 2     | Diclofenac sod. | 10              | 0.10±0.015* | 0.13±0.006* | 0.09±0.00** | 0.06±0.001** | 71.31        |
| 3     | EENZ            | 250             | 0.21±0.012  | 0.24±0.012  | 0.19±0.008  | 0.15±0.010*  | 39.22        |
| 4     | EENZ            | 500             | 0.13±0.018* | 0.18±0.014* | 0.13±0.008* | 0.09±0.002** | 59.23        |

Values are mean ± SEM (n = 6). \* $p < 0.05$ , \*\* $p < 0.001$  when compared to control

**Table 3. Anti-inflammatory effect of EENZ on Serotonin induced rat paw edema**

| Sr no | Treatment       | Dose mg/kg p.o. | 0.5h       | 1h          | 3h           | 5h           | %inhibition |
|-------|-----------------|-----------------|------------|-------------|--------------|--------------|-------------|
| 1     | Control         |                 | 0.26±0.02  | 0.29±0.015  | 0.35±0.020   | 0.38±0.011   | -           |
| 2     | Diclofenac sod. | 10              | 0.18±0.018 | 0.15±0.015* | 0.10±0.014** | 0.08±0.001** | 63.42       |
| 3     | EENZ            | 250             | 0.24±0.020 | 0.22±0.005  | 0.19±0.013*  | 0.16±0.020*  | 38.31       |
| 4     | EENZ            | 500             | 0.20±0.016 | 0.17±0.014  | 0.13±0.012*  | 0.11±0.011** | 54.19       |

Values are mean ± SEM (n = 6). \* $p < 0.05$ , \*\* $p < 0.001$  when compared to control

**Table 4. Anti-inflammatory effect of EENZ on Cotton pellet induced granuloma**

| Treatment       | Dose mg/kg | Mean Wet Weight of Pellet (mg) | Percentage inhibition | Mean Dry Weight of Pellet (mg) | Percentage inhibition |
|-----------------|------------|--------------------------------|-----------------------|--------------------------------|-----------------------|
| Control         |            | 198.68 ±10.47                  | -                     | 45.60 ±2.04                    | -                     |
| Diclofenac sod. | 10         | 95.30 ±3.46 *                  | 52.03                 | 20.88 ±0.77 *                  | 54.21                 |
| EENZ            | 250        | 157.68 ±8.18                   | 20.64                 | 37.40 ±1.59                    | 17.98                 |
| EENZ            | 500        | 115.15 ±7.43 *                 | 42.43                 | 25.63 ±1.53 *                  | 44.26                 |

Values are mean ± SEM (n = 6). \* $p < 0.05$  when compared to control

Source of support: Nil, Conflict of interest: None Declared