



ANTIBACTERIAL ACTIVITY OF A NEW FLAVONE GLYCOSIDE FROM THE SEEDS OF *CASSIA SOPHERA* LINN.

Jain Parul^{*1} and Nema Rajeev²

¹Department of Chemistry, S. V. Poly College Bhopal, M.P., India

²S.N.G.G.P.G. Collage Department of Zoology Bhopal, M.P., India

Article Received on: 20/01/12 Revised on: 11/03/12 Approved for publication: 25/03/12

*Email: drparuljainsv@gmail.com

ABSTRACT

A new flavone glycoside 1, m.p. 248-251°C, m.f. C₃₃H₄₀O₂₀, [M]⁺ 756 (FABMS) has been isolated from methanol soluble fraction of defatted seeds of *Cassia sophera* Linn., which was characterized as 5,7,3',4'-tetrahydroxy-3-methoxyflavone-5-O-α-L-rhamnopyranosyl-7-O-β-D-glucopyranosyl (1→3)-O-β-D-xylopyranoside by various colour reactions, chemical degradations and spectral analyses. In present paper, it was screened against *Bacillus coagulans*, *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa* by using paper disc diffusion method. Results were compared with the zone of inhibition produced by commercially available standard antibiotic. It was observed that it showed good activity against these microbes.

KEYWORDS: antibacterial activity, flavone glycoside, *Cassia sophera* Linn.

INTRODUCTION

The natural products play an important role against microorganisms. Plants are unlimited source of natural products. They still form a major part of ingredient in almost all system of therapeutics. There are more than 350 thousand species of higher plants out of which a small portion has been investigated for their biologically active constituents.

The biological screening of indigenous drugs from plants was started in India by Col R.N. Chopra¹. Kurup² studied the presence of antibacterial agent in Ayurvedic system of medicine. The antimicrobial activity of extracts of leaves of *Spondias mombin* and *Alchornea cordifolia* have been carried out by Ajao and his coworker³. Among the numerous products identified from medicinal plants, flavonoids represent one of the most interesting groups of biologically active compounds. Flavonoids are polyphenolic compounds isolated from wide varieties of vascular plants with over 8000 individual compounds known⁴. Flavonoids are plant pigments which occur in the plant kingdom and occur naturally in free state (aglycone), or as glycosides, or associated with tannins, possessing secretory structures. The most common classes are flavonol, flavones and their dihydroderivatives followed by anthocyanins, flavans and isoflavans. Flavonoids are found in vegetables, fruits, seeds, nuts, grains, spices and different medicinal plants as well as in specific beverages such as red wine, tea and unfiltered beer⁵. They are important constituents of the non energetic parts of the human diet, the average intake being around of 600 mg/day⁶. Protective role of flavonoid intake against coronary heart diseases has been reported in epidemiological studies⁷. Several flavonoids have been considered valuable phytochemicals for different body system: urinary, digestive, cardiovascular, nervous and skin⁸. Various flavonoids isolated from the plants have been shown to antimicrobial⁹⁻¹⁰, antiviral¹¹, antiinflammatory¹², antimutagenic¹³⁻¹⁴, anticancer¹⁵, antiulcerogenic¹⁶, antioxidant¹⁷ and antitumor¹⁸ activity. *Cassia sophera* Linn.¹⁹⁻²⁰ (Leguminosae) is commonly known as 'Kasundi' or 'Banar' in Hindi. It is a shrub or under shrub 2.4-3 m high, annual or perennial. It is distributed throughout India and in most tropical countries. Its bark and seeds are useful in diabetes.

Its leaves are used externally in ringworm. Decoction of plant is used in acute bronchitis. Its bark, leaves and seeds are used as cathartic. We have already reported the isolation and structural elucidation of a new flavone glycoside²¹ from the seeds of this plant. The methanol fraction of the defatted seeds of the plant afforded a new flavones glycoside(1), m.p. 248-251°C, C₃₃H₄₀O₂₀ and [M]⁺ 756 (FABMS), which was characterised as 5,7, 3', 4'-tetrahydroxy-3-methoxyflavone-5-O-α-L-rhamnopyranosyl-7-O-β-D-glucopyranosyl(1→3)-O-β-D-xylopyranoside by various colour reactions chemical degradations and spectral analyses²¹. In the present paper, we report the antibacterial activity of this new flavone glycoside (1).

MATERIAL AND METHODS

Plant Material

The seeds of *Cassia sophera* were procured from M/S United Chemicals and Allied Products Kolkata and were taxonomically authenticated by the Department of Botany, Dr. H.S. Gour University Sagar (M.P.), India. A voucher specimen has been deposited in the Natural Products Laboratory, Department of Chemistry, Dr. H.S. Gour University, Sagar (M.P.), India.

Extraction and Isolation

Air-dried powdered seeds (3 kg) of the plant were extracted with petroleum ether (40-60°C) in a Soxhlet apparatus for seven days. The defatted seeds successively extracted with C₆H₆, CHCl₃, CH₃COOCH₃, CH₃COCH₃ and methanol. The methanol soluble fraction of the plants was concentrated under reduced pressure which showed two spots on TLC examination using solvent system CHCl₃:MeOH (4:6) indicating it to be a mixture of 1 and 2. These compounds were separated and by column chromatography over silica gel and purified by preprative TLC and studied separately. Compound 1 was characterised as 5,7,3',4'-tetrahydroxy-3-methoxyflavone-5-O-α-L-rhamnopyranosyl-7-O-β-D-glucopyranosyl (1→3)-O-β-D-xylopyranoside by various colour reactions chemical degradations and spectral analyses²².

Test microorganism

The test microorganism used were: *Bacillus coagulans*, *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas*

aerugenosa. Paper disc agar diffusion method²² were used for in vitro evaluation of antibacterial activity.

Preparation of culture media

Peptone-beef extract has been used for study of antibacterial activity with the following composition:

Peptone	5.0 gm
Beef extract	3.0 gm
Sodium chloride	5.0 gm
Agar	15.0 gm
Distilled water	1000 ml

All the above ingredients were dissolved in distilled water and volume made up to 1 litre. The pH of the medium was adjusted to 7.2 with 1N NaOH.

Preparation of slants

5 ml of sterilized nutrient agar was poured in sterilized culture tube and allowed to cool. The tubes were incubated in electrically heated incubators at different temperature for various periods for different kinds of microorganisms depending upon the optimum growth of microorganisms.

Sterilization

The media and the slants were sterilized by autoclaving for about 15-20 minutes at 15 lb pressure for the preparation of

subcultures of the organism. Petridishes were sterilized in an electrically heated oven at 125°C for 8 hrs.

Preparation of agar plates

The sterilized media was cooled to 45°C. Homogeneous suspension of each microorganism was mixed with sterilized media and 15 ml of this medium was poured into each sterilized Petridish and allowed to gel.

Standard drug used

Streptomycin was used as standard antibacterial agent.

Antibacterial activity

Petri plates were pre-seeded with 15 ml of growth agar medium and 1.0 ml of inoculum²²⁻²³. Paper discs of 6 mm diameter, which absorbs about 0.1 ml of test samples of the compound **1** and known quantity of standard reference antibiotic, were used. The inoculated plates were kept at 5 °C for 40-45 minutes so as to allow the diffusion of the substances and then incubated at 36±1 °C for 36 hours. The inhibition zones were measured in mm and the results obtained are recorded in the **Table-I** and compared with the standard reference antibiotic²⁴. On the basis of **Table-I**, various graphs has been drawn to show the comparison of antibacterial activity between compound and standard drug against each single bacteria.

TABLE-I ANTIBACTERIAL ACTIVITY OF COMPOUNDS

Compound	Diameter of Zone of inhibition (mm) against															
	(+) <i>Bacillus coagulans</i>				(+) <i>Staphylococcus aureus</i>				(-) <i>Escherichia coli</i>				(-) <i>Pseudomonas aeruginosa</i>			
	Concentration of compound				Concentration of compound				Concentration of compound				Concentration of compound			
	100%	80%	60%	40%	100%	80%	60%	40%	100%	80%	60%	40%	100%	80%	60%	40%
Pa	17.2	14.3	10.1	3.5	18.3	15.0	12.7	9.4	11.1	8.3	4.8	3.0	9.8	7.0	3.1	-
**Std.	21.2	20.0	19.5	18.0	20.0	19.1	18.3	17.9	18.3	17.0	15.6	14.0	23.1	21.0	19.5	17.0

*The zone of inhibition (mm) taken as average of four determinations in four different directions.

**Streptomycin used as standard antibacterial agent.

RESULTS AND DISCUSSION




Fig 1. Screening activity of compound 1 with the spatial reference to streptomycin against *Bacillus coagulans*




Fig 2 Screening activity of compound 1 with the spatial reference to streptomycin against *Staphylococcus aureus*




Fig 3. Screening activity of compound 1 with the spatial reference to and streptomycin against *Escherichia coli*.




Fig 4. Screening activity of compound 1 with the spatial reference to streptomycin against *Pseudomonas aeruginosa*

The results reported in **Table-1** showed that the antibacterial activity of compound **1**, was found to be fairly good against gram positive bacteria *Staphylococcus aureus* and *Bacillus coagulans*. It also exhibited good activity against gram negative bacteria *Escherichia coli* and it showed significant activity against gram negative bacteria *Pseudomonas aeruginosa* only on higher concentration. Thus on the basis of above results, it is obvious that the above flavonoid glycoside (**1**) may be potentially used as antibacterial agent.

ACKNOWLEDGMENT

Authors are grateful to Prof. R. N. Yadava, Department of Chemistry, Dr. H.S. Gour University, Sagar (M.P.), for his support, suggestion and encouragement throughout the production of paper. We also want to acknowledge to the Head, Department of Botany of this University for providing antimicrobial facilities.

REFERENCES

- Chopra RN, Handa IC and Kapur LD. Indigenous Drugs of India, U.N. Shart and Sons, Calcutta, 2nd ed. 1958; 422.
- Kurup PA. Studies on plant antibiotics: Screening of some Indian medicinal plants. J. Sci. Ind. Res. 1956; 15C: 153-154.
- Ajao AO, Shonukan O and Femi-Onadeko B. Antibacterial effect of aqueous and alcohol extracts of *Spondian mombin* and *Alchronea cordifolia*: Two local antimicrobial remedies. Int. J. Crude Drugs Res. 1985; 23: 67.
- Harborne, JB. The Flavonoids: Advances in Research, Chapman and Hall, London 1994.
- Kuhnau, J. The flavonoids; a class of semi essential food components; their role in human nutrition World Rev. Nutr. Diet. 1976; 24: 117-191.
- Manach, C, Regerat F, Texier O, Agullo G, Demigne C, Remesy C. Bioavailability, metabolism and physiological impact of 4-oxo-flavonoids. Nutr. Res. 1996; 16: 517-544.
- Hertog MGL and Katan MB. The Flavonoids in Health and Disease, Rice-Evans, C. A., Packer, L. Eds. Marcel Dekker, New York 1998; 447-467..
- Tyler VE. The Therapeutic use of Phytomedicinals, Haworth Press, Inc., New York 1995.
- Harsh ML, Nag TN. and Jain S. Arid zone plants of Rajasthan-A source of antimicrobials. ; Comp. Physiol.Ecol. 1983; 2: 129-131.
- Jit S, Shekhawat SS, Grover S, Nag TN. Screening of some plant of Zygophyllaceae for their antimicrobial activity. Acta Botanica. Indica. 1986; 14: 45-47.
- Mahmood N, Pizsa C, Aquino R, De Tommasi N, Piacente S, Colman S, Burke A. and Hay A. nhibition of HIV infection by flavonoids. J. Antiviral. Res.1993; 22: 189-99.
- Carvalho, J.C.T., Ferreira, L.P.,Dasilva Santos, L., Correa, M.J.C., Oliveira. Campos, L.M., Bastos, J.K. and Sarti, S.J.; Antiinflammatory activity of flavone and some of its derivates from *virola michelli* Heckel. J. Ethanopharmacol. 1999 ;64(2): 173-177.
- Brit, D.F., Walker, B., Tibbles, M.G. and Bresnic, E.; Anti-mutagenesis and antipromotion by apigenin robinetin and indole-3-carbinol. Carcinogenesis 1986; 7(6): 959-963.
- Wall ME, Wani MC, Hughes TJ. and Taylor H. Plant antimutagenic agents, I. General bioassay and Isolation procedures. J. Nat. Prod.1986; 51: 866-873.
- Nixon JE, Hendricks JD, Parolowski NE, Perrira,CB, Sinhuber RO. and Bailey GS. Inhibition of aflatoxin B1 carinogenesis in rainbow trout by flavones and indole compounds. Carcinogenesis1984; 5: 615-619.
- Kaneda, N, Pezzuto JM, Soefarto DD, Kingnorn AD. and Fransworth NR. Plant anticancer agents,XLVII. New cytotoxic flavonoids from *Mutingia Colabura* roots. J. Nat. Prod. 1991; 54(1): 196-206.
- Gamez EJC, Luyengi L, Lee SK, Zhu LF, Zhou BN, Fong HHS, Joh JM, Pezzuto JM. and Kingnorn AD. Antioxidant flavonoid glycosides from *Daphniphyllum Calycinum*. J. Nat. Prod., 1998 ;61: 706-708.
- Ulubelen A, Bucker R. and Mabry TJ. Flavone-5-O-glycoside from *Daphne seri-cea*. Phytochemistry 1982; 21: 801-803.
- Chopra RN, Nayer SL. and Chopra IC. Glossary of Indian Medicinal Plants, (CSIR Publication, New Delhi) 1956; 55.
- Kritikar KR. and Basu BD. Indian Medicinal Plants, 2nd edition, Lalit Mohan Basu and Co., Allahabad 1948; Vol. II: 863-865.
- Yadva RN. and Jain P. A new flavone glycoside from *Cassia sophera* Linn. J. Indian Chem. Soc. 2006; 83: 1175-1178.
- Khanna P, Mohan S. and Nag TN. Antimicrobials from plant tissue cultures. Lloydia.1971; 34: 168-169.
- Harsh ML. and Nag TN. Antimicrobial principles from in vitro tissue culture. Lloydia. 1984; 47: 365-368

Source of support: Nil, Conflict of interest: None Declared