

SYNTHESIS AND BIOLOGICAL EVALUATION OF SOME PHENYL ACETIC ACID HYDRAZONE DERIVATIVES

Jadon Gunjan^{*1}, Diwaker Awadh²¹Department of Pharmaceutical Chemistry, Shrinathji Institute of Pharmacy, Nathdwara, dist. Rajsamand (RAJ.)²Department of Biotechnology, JSS College of pharmacy, Ooty, India

Article Received on: 05/09/11 Revised on: 22/10/11 Approved for publication: 09/11/11

*E-mail: Jadon_gunjan@yahoo.in

ABSTRACT

These products are the result of further operation after the derivation of three hydrazones derivative compounds-

N'-(4-Chlorobenzylidene)-2-phenylacetohydrazide

N'-(2-chlorobenzylidene)-2-phenylacetohydrazide

N'-(3, 4, 5-trimethoxybenzylidene)-2-phenylacetohydrazide

N'-(furan-2-ylmethylene)-2-phenylacetohydrazide

N'-(3, 4, 5-Trimethoxybenzylidene)-2-phenylacetohydrazide

N'-(1-(4-hydroxyphenyl) ethylidene)-2-phenylacetohydrazide

These make a considerable interest in process development of novel compounds with anticonvulsant, antidepressant, analgesic, and anti-inflammatory, anti platelet, anti malarial, antimicrobial, antimycobacterial, antitumoral, vasodilator, and antiviral and anti chistosomiasis activities. These all three compounds evaluated their biological activities. These observations have been guiding for the development of new hydrazones that possess varied biological activities.

Keywords: Hydrazones, Hydrazide-Hydrazones, Phenyl Acetic Acid, Anti-inflammatory Activity, Biological Activity.


INTRODUCTION

Hydrazones have been demonstrated to possess, among other, antimicrobial, anticonvulsant, and analgesic, anti-inflammatory, and ant platelet, ant tubercular and antitumoral activities. For example, isonicotinoyl hydrazones are antitubercular; 4-hydroxybenzoic acid[(5-nitro-2-furyl)ethylene]-hydrazide (nifuroxazide) is an intestinal antiseptic; 4-fluorobenzoic acid[(5-nitro-2-furyl)methylene]-hydrazide¹ and 2,3,4-pentanetrione-3-[4-[[[(5-nitro-2-furyl)methylene]hydrazino]carbonyl]phenyl]-hydrazone², which were synthesized in our Department, have antibacterial activity against both *Staphylococcus aureus* ATCC 29213 and *Mycobacterium tuberculosis* H37Rv at a concentration of 3.13 µg/mL. N1-(4-Methoxybenzamido)benzoyl]-N2-[(5-nitro-2-furyl)methylene]hydrazine, which was also synthesized in our Department³, demonstrated antibacterial activity. In addition, some

of the new hydrazide-hydrazones that we have recently synthesized were active against the same strain of *M. tuberculosis* H37Rv between the concentrations of 0.78-6.25 µg/mL⁴.

General procedure

Hydrazide-hydrazones compounds are not only intermediates but they are also very effective organic compounds in their own right. When they are used as intermediates, coupling products can be synthesized by using the active hydrogen component of –CONHN=CH– azomethine group⁸. N-Alkyl hydrazides can be synthesized by reduction of hydrazones with NaBH₄⁹, substituted 1,3,4- oxadiazolines can be synthesized when hydrazones are heated in the presence of acetic anhydride^{1,10,11}. 2-Azetidinones can be synthesized when hydrazones react with triethylamine chloro acetylchloride¹². 4-Thiazolidinones are synthesized when hydrazones react with thioglycolic acid/ thiolactic acid^{3,13}.



Scheme -1

OUTLINE ABOUT PREVIOUS COMPOUNDS

Comp.Code	Name	Mol. Formula	Mol. Wt	M.P. (°C)
GL ¹	N'-(4-Chlorobenzylidene)-2-phenylacetohydrazide	C ₁₅ H ₁₃ N ₂ OCl	272	61
GL ²	N'-(2-Chlorobenzylidene)-2-phenylacetohydrazide	C ₁₅ H ₁₃ N ₂ OCl	272	60.5
GL ³	2-Phenyl-N'-(1-(2,3,4-trimethoxyphenyl)ethylidene)acetohydrazide	C ₁₃ H ₁₂ N ₂ O ₂	228	55.5
GL ⁴	N'-(furan-2-ylmethylene)-2-phenylacetohydrazide	C ₁₈ H ₂₀ N ₂ O ₄	328	50.5
GL ⁵	N'-(3,4,5-Trimethoxybenzylidene)-2-phenylacetohydrazide	C ₁₇ H ₁₇ N ₂ O ₂	281	71
GL ⁶	N'-(1-(4-hydroxyphenyl)ethylidene)-2-phenylacetohydrazide	C ₁₆ H ₁₆ N ₂ O ₂	268	51


MATERIALS AND METHODS

All the melting points reported in this dissertation progress report were determined by open capillary tube method and are uncorrected. The synthetic and analytical studies of the compounds were carried out using laboratory grade and analytical grade reagents as the case may be standard procedure or reported methods were followed with or without modification appropriately as and when required.

General methodology for experimental compound

Synthesis of N'-(1-(2-hydroxyphenyl)ethylidene)-2-phenylacetohydrazide: (GL⁷)


A mixture of o-hydroxy acetophenone (1.36gm, 0.01mole) and 2-phenyl acetohydrazide (1.5gm, 0.01mole) were dissolved in methanol then two drops of conc. HCl were added as catalyst and stirred at room temperature for 5 hr. the reaction mixture was poured into ice and filtered. The crude product so obtained was dried and recrystallized with methanol.



N'-(1-(2-hydroxyphenyl)ethylidene)-2-phenylacetohydrazide

Synthesis of N'-(1-phenyl ethylidene)-2-phenylacetohydrazide: (GL⁸)

A mixture of acetophenone (1.28gm, 0.01mole) and 2-phenyl acetohydrazide (1.5gm, 0.01mole) were dissolved in methanol then two drops of conc. HCl were added as catalyst and stirred at room temperature for 4 hr. the reaction mixture was poured into ice and filtered. The crude product so obtained was dried and recrystallized with methanol.



General methodology Scheme-2

Table - 1: The Physico-chemical data of synthesized compounds

Comp.Code	Mol. Formula	Mol. Wt.	F _{ro}	Color (Appearance)	M.P. (°C)	λ _{max}
GL ⁷	C ₁₆ H ₁₆ N ₂ O ₂	268	0.85 ^b	Black	53	258.329
GL ⁸	C ₁₆ H ₁₆ N ₂ O	252	0.90 ^c	Light brown	61.5	330.805

Solvent System:

a. (Chloroform: Methanol 8:2), b. (Chloroform: Methanol 8 : 1)

Table -2: IR Data of the title compounds

Compound	IR (cm ⁻¹)
GL ⁷	3396.67(OH), 3157.1(NH), 2951.76(Ar-CH), 2896(CH ₃) ₃ , 1685.3(C=O), 1630.54(C=N)

BIOLOGICAL ACTIVITY OF HYDRAZONE COMPOUNDS

Anticonvulsant Activity

Epilepsy is a common neurological disorder and a collective term given to a group of syndromes that involve spontaneous, intermittent, abnormal electrical activity in the brain. The pharmacotherapy of epilepsy has been archived during the last decade. Furthermore, although for the last twenty years new antiepileptic drugs have been introduced into clinical practice, the maximal electroshock (MES) test and the subcutaneous pentylenetetrazole (scPTZ) test are the most widely used animal models of Epilepsy to characterize the anticonvulsant activity. The biological results revealed that in general, the acetylhydrazones **2** provided good protection against convulsions while the oxamoylhydrazones **3** were significantly less active.²¹

Analgesic, Anti-inflammatory and Anti platelet Activity

Non-steroidal anti-inflammatory drugs (NSAIDs) have a wide clinical use for the treatment of inflammatory and painful conditions including rheumatoid arthritis, soft tissue and oral cavity lesions, respiratory tract infections and fever. The two isoforms of cyclooxygenase (COX) are poorly distinguishable by most of the classical NSAIDs and these agents actually inhibit COX-1 extensively, besides COX-2, leading to gastrointestinal injury,

suppression of TXA₂ formation and platelet aggregation. The combination of these interactions is probably the reason for gastrointestinal bleeding as the most serious complication of these drugs. Some evidences suggest that the hydrazone moiety present in some compounds possess a pharmacophoric character for the inhibition of COX. The most important anti-inflammatory derivative 2-(2-formylfuryl) pyridylhydrazone **7** presented a 79 % inhibition of pleurisy at a dose of 80.1 mol/kg. The authors also described the results concerning the mechanism of the action of these series of N-heterocyclic derivatives in platelet aggregation that suggests a Ca²⁺ scavenger mechanism. Compound **7** was able to complex Ca²⁺ in invitro experiments at 100 concentrations, indicating that these series of compounds can act as Ca²⁺ scavenger depending on the nature of the aryl moiety present at the imine subunit.²²

Antimycobacterial Activity

Tuberculosis is a serious health problem that causes the death of some three million people every year worldwide²³. In addition to this, the increase in *M. tuberculosis* strains resistant to front-line antimycobacterial drugs such as rifampin and INH has further complicated the problem, which clearly indicates the need for more effective drugs for the efficient management of tuberculosis. Meyer

and Mally prepared new hydrazones by reacting isoniazid (INH) with benzaldehyde, o-chlorobenzaldehyde and vanillin⁵. Shchukina *et al.* prepared INH hydrazone-hydrazones **1** by reacting INH with various aldehydes and ketones; the compounds were reported to have activity in mice which had been infected with various strains of *M. tuberculosis*, and also indicated lower toxicity than INH^{5,6}.

The reaction of 1-methyl-1H-2-imidazo [4, 5-b] pyridine carboxylic acid hydrazide with substituted aldehydes yielded the corresponding hydrazone-hydrazones. Compound **32** exhibited antimycobacterial activity against *M. tuberculosis* H37 Rv, *M. tuberculosis* 192, *M. tuberculosis* 210, isolated from patients and resistant against INH, ethambutol, rifampicine at 31.2 ig/mL²⁴.

Antitumoral Activity

A variety of antitumoral drugs are currently in clinical use. The search for antitumoral drugs led to the discovery of several hydrazones having antitumoral activity. Some of diphenolic hydrazones showed maximum uterotrophic inhibition of 70%, whereas compound **58** exhibited cytotoxicity in the range of 50-70% against MCF-7 and ZR-75-1 human malignant breast cell lines²⁵.

Antimicrobial Activity

The dramatically rising prevalence of multi-drug resistant microbial infections in the past few decades has become a serious health care problem. The search for new antimicrobial agents will consequently always remain as an important and challenging task for medicinal chemists. Ethyl 2-arylhydrazono-3-oxobutyrate **17** were synthesized in order to determine their antimicrobial properties. Compound **17d** showed significant activity against *S. aureus* whereas the others had no remarkable activity on this strain. Compound **17e** was found to be more active than the others against *Mycobacterium fortuitum* at a MIC value of 32 ig/ml²⁶.

RESULT & DISCUSSION

The synthesized compounds were purified and the purity of all compounds was checked by melting point and RF value determination. The structures of this compound were established by recording spectral data (Table-1, 2 & 3). In the present work synthesized hydrazones derivative compound significantly expected to show Anticonvulsant Activity, Analgesic, Anti-inflammatory Antiplatelet Activity Antimycobacterial Activity, and Antimicrobial Activity.

REFERENCES

1. Rolla's S, Gülerman N, Erdeniz H. Synthesis and antimicrobial activity of some new Hydrazones of 4-fluorobenzoic acid hydrazide and 3-acetyl-2, 5-disubstituted-1, 3, 4-oxadiazolines. *Farmaco* 2002; 57: 171-174
2. Kuçukguzel G, Rolla's S, Kucukguzel y, Kara M. Synthesis and Antimycobacterial activity of some coupling products from 4-aminobenzoic acid hydrazones. *Euro. J. Med. Chem.* 1999; 34: 1093-1100
3. Kucukguzel G, Oruc E E, Rolla's S, akin F, ozbek A. Synthesis, Characterization and biological activity of novel 4-thiazolidinones, 1,3,4-oxadiazoles and some related compounds. *Euro. J. Med. Chem.* 2002; 37: 197-206
4. Kaymakçoolu K B, Oruc E E, Unsalan S, Kandemirli F, Shvets N, Rollas S, Anatholy D. Synthesis and characterization of novel hydrazide-hydrazones and the study of their structureantituberculosis activity. *Euro. J. Med. Chem.* 2006; 41: 1253-1261

5. Sah P P T, Peoples S A. Isonicotinyl hydrazones as antitubercular agents and derivatives for identification of aldehydes and ketones. *J. Am. Pharm. Assoc.* 1954; 43: 513-524
6. Bevin E M, Drain D J, Seiler M, Seymour D E. Some further studies on tuberculostatic compounds. *J. Pharm. Pharmacology.* 1954; 4: 844-855
7. Buu-Hoi P H, Xuong D, Nam H, Binon F, Royer R. Tuberculostatic hydrazides and their derivatives. *J. Chem. Soc.* 1953; 1358-1364
8. Singh V, Srivastava V K, Palit G, Shanker K, Coumarin congeners as antidepressants. *Arzneim-Forsch. Drug. Res.* 1992; 42: 993-996
9. Ergenc N, Gunay N S. Synthesis and antidepressant evaluation of new 3-phenyl-5-sulfonamidoindole derivatives. *Eur. J. Med. Chem.* 1998; 33: 143-148
10. Durgun B B, capan G, Ergenç N, Rollas S. Synthesis, characterisation and biological evaluation of new benzyldenebenzohydrazides and 2,5-disubstituted-2,3-dihydro-1,3,4-oxadiazoles. *Pharmazie* 1993; 48: 942-943
11. Dooan H N, Duran A, Rolla's S, Ener G, Armutak Y, Keyer-Uysal M. Synthesis and structure elucidation of some new hydrazones and oxadiazolines anticonvulsant activities of 2-(3-acetyloxy-2-naphthyl)-4-acetyl-5-substituted-1,3,4-oxadiazolines. *Med. Sci. Res.* 1998; 26: 755-758
12. Kalsi R, Shrimali M, Bhalla T N, Barthwal J P. Synthesis and anti-inflammatory activity of indolyl azetidinones. *Indian J. Pharm. Sci.* 2006; 41: 353-359
13. Kucukguzel G, Kocatepe A, De Clercq E, ahin F, Gulluce M. Synthesis and biological activity of 4-thiazolidinones, thiosemicarbazides derived from diflunisal hydrazide. *Euro. J. Med. Chem.* 2006; 41: 353-359
14. Komurcu G, Rollas S, Ulgen M, Gorrod J W, Cevikba A. Evaluation of Some Arylhydrazones of p-Aminobenzoic acid hydrazide as Antimicrobial Agents and Their in-vitro Hepatic Microsomal Metabolism. *Boll. Chim. Farm.* 1995; 134: 375-379
15. Ulgen M, Durgun, B B, Rolla's S, Gorrod J W. The in vitro hepatic microsomal metabolism of benzoic acid benzyldenehydrazide. *Drug Metab. Interact.* 1997; 13: 285-294
16. Vora J J. *et al.* 2009; have reported synthesis and microbial studies of some novel Schiff bases derivatives of 4-methyl pyridine-2-amine
17. Rajput A P. *et al.* 2009; Have synthesized benzaldehyde substituted phenyl carbonyl hydrazones and their formyl derivatives using vilsmeier-Haack-reaction
18. Sham M Sondhi. *et al.* 2009; Have reported synthesis of biologically active novel bi's schiff bases and bi's hydrazones derivatives
19. Sandhya B. *et al.* 2008; Have synthesized synthesis of Schiff's bases of 8-methyl - Tetrazolo (1, 5-a) quinoline as potential anti- inflammatory and antimicrobial agents
20. Olcay Bekiracan. *et al.* 2008; Have reported synthesis of Schiff's and mannich bases Of isatin derivatives with 4-amino-4, 5-dihydro-1H-1, 2, 4-triazole-5-ones
21. Dimmock J R, Vashishtha S C, Stables J P. Anticonvulsant properties of various Acetyl hydrazones, oxamoylhydrazones and semicarbazones derived from aromatic and
- Unsaturated carbonyl compounds. *Euro. J. Med. Chem.* 2000; 35: 241-248
22. Todeschini A R, Miranda A L, Silva C M, Parrini S C, Barreiro E J. Synthesis And evaluation of analgesic, anti-inflammatory and ant platelet properties of new 2-Pyridylarylhydrazone derivatives. *Euro. J. Med. Chem.* 1998; 33: 189-199
23. www.TAACF.org.
24. Bukowski L, Janowicz M. 1-Methyl-1H-2-imidazo [4, 5-b] pyridine carboxylic acid and some of derivatives with suspected antituberculous activity. *Pharmazie* 1996; 51: 27-30
25. Pandey J, Pal R, Dwivedi A, Hajela K. Synthesis of some new diaryl and triaryl hydrazones derivatives as possible estrogen receptor modulators. *Arzneimittelforschung.* 2002; 52: 39-44
26. Kucukguzel G, Rolla's S, Erdeniz H, Kiraz M. Synthesis, Characterization and Antimicrobial Evaluation of Ethyl 2-Arylhydrazono-3-oxobutyrate, *Euro. J. Med. Chem.* 1999; 34: 153-160
27. Walcourt A, Loyevsky M, Lovejoy D B, Gordeuk V R, Richardson D R. Novel Aroylhydrazone and thiosemicarbazone iron chelators with anti-malarial activity against Chloroquine-resistant and -sensitive parasites. *Int. J. Biochem. Cell Biol.* 2004; 36: 401-407

Source of support: Nil, Conflict of interest: None Declared