

Research Article



INTERNATIONAL RESEARCH JOURNAL OF PHARMACY
www.irjponline.com
ISSN 2230-8407 [LINKING]

Morphometric analysis of the proximal end of a desiccated human femur and its clinical implications.

Dr.Asha Changeli¹, Dr.Anjali Patil², Dr.Rajendra singh³

1. Ph.D Scholar, Dept of Anatomy Malwanchal University, Indore(dr.ashi09@gmail.com)
2. Associate Professor, Dept.of Anatomy, IMCH&RC, Indore
3. Assistant Professor, Dept.of Anatomy, IMCH&RC, Indore(rajendrasingh211@gmail.com)

Corresponding Author: Dr.Asha Changeli,email: dr.ashi09@gmail.com

Abstract:

Background: The longest and strongest bone in the human body, the femur, is essential for weight transfer and bipedal movement. Its size and form vary widely according on the environment, gender, and ethnicity. **Aim:** In order to comprehend the clinical consequences in orthopedic treatment, the study intends to investigate the morphometric characteristics of dehydrated human femora, including the head, neck, and trochanters. The information will be used to forecast fracture risk, implant fit, and surgical results, as well as to help design and choose orthopedic implants, especially for hip joint replacement procedures. **Materials & methods:** A total of 50 dry human femora were included in the study. Only complete, undamaged adult femora with clearly identifiable anatomical landmarks were selected. Bones with signs of deformity, pathology, or damage at the proximal end were excluded from the analysis. Femur measures such as neck diameter, length, thickness, shaft angle, intertrochanteric line length, maximum femur length, vertical head diameter, foveal diameter, fovea transverse diameter, and foveal longitudinal diameter were examined in the study. Measurements were taken using the following tools: Digital Vernier Caliper (accuracy: 0.01 mm), Measuring tape, Osteometric board, and Goniometer. **Results:** Femur measures such as neck diameter, length, thickness, shaft angle, intertrochanteric line length, and maximum femur length were examined in the study. 29.47 ± 3.43 , 38.06 ± 4.04 , 29.61 ± 1.93 , 121.28 ± 5.38 , 43.14 ± 3.22 , and 44.11 ± 2.91 were the mean values. In these parameters, there was no discernible variation between gender and age. Measurements of the femur, such as the vertical head diameter, foveal diameter, fovea transverse diameter, and foveal longitudinal diameter, were also examined in the study. The mean foveal depth was 2.87 ± 1.24 mm, and the vertical head diameter was 41.51 ± 4.27 mm. The longitudinal and transverse dimensions of the fovea were 17.16 ± 5.37 mm and 13.58 ± 1.97 mm, respectively. There were no discernible gender or age disparities. **Conclusion:** In order to guide implant design and selection, this study examined morphometric characteristics in dehydrated human femora and demonstrated their importance in orthopedic health and surgical procedures.

Key words: Neck diameter, neck length, neck thickness, neck shaft angle, maximum length of the femur, foveal diameter, and fovea transverse diameter.

Introduction

Bipedal movement and the transfer of weight from the axial skeleton to the lower extremities depend on the human femur, the longest and strongest bone in the body. The environment, gender, place of residence, and ethnic heritage may all have a significant impact on the size and form of the femur's upper portion, which comprises the femoral head, neck, greater trochanter, and lesser trochanter. Understanding these differences is essential for anthropological research, prosthetic design, forensic identification, and orthopedic procedures.

Numerous morphometric investigations have been conducted on the proximal femur in India, revealing regional and ethnic differences [1]. According to a research [2], femoral head diameter and neck-shaft angles differed from Western norms, highlighting the necessity for region-specific data. Similarly, studies in central India [3] found significant differences in the size of the proximal femur between boys and girls. This emphasizes how important gender-specific anatomical databases are for therapeutic use.

Globally, morphometric investigations have revealed notable differences amongst populations. According to research from Europe, Africa, and East Asia, the observable disparities in femur sizes among races have a significant impact on the design of orthopedic implants and forensic estimations. For example, a study [4] showed a wider femoral head and a greater neck-shaft angle compared to Caucasians, which may affect hip arthroplasty outcomes. Neck lengths were shorter and femoral head diameters were smaller, according to studies like those by [4-6]. Custom prosthetic models were therefore necessary to provide the best possible fit and performance.

Furthermore, comparative studies have emphasized that information gathered from Western populations cannot be used directly to Indian patients without running the risk of surgical problems and implant mismatch. According to a comparative investigation, Indian patients often experience worse than ideal results from traditional prosthetic designs that are based on Western anthropometry [7]. They advocated for the use of domestic data-driven implant design methodologies. Furthermore, clinical consequences such as fracture risk assessment, surgical planning for total hip replacements (THR), and exact forensic profiling depend on precise morphometric information of the proximal femur [2,3]. The morphometry of dehydrated femora is particularly important in forensic anthropology and bioarchaeology since dried bone specimens are commonly used as the main source material for studies. The measurements taken from dried bones are crucial for determining age and sex, reconstructing stature, and identifying unknown people. Forensic investigations have benefited from the formulae developed by Indian research, such as those by [7,8], to estimate height using measurements taken from the top section of the thigh bone. Meanwhile, worldwide studies, including the one by [10–12], have improved sex determination techniques based on femoral head measures, highlighting the diagnostic utility of the bone. In order to comprehend the clinical consequences in orthopedic treatment, the study intends to investigate the morphometric characteristics of dehydrated human femora, including the head, neck, and trochanters. The information will be used to forecast fracture risk, implant fit, and

surgical results, as well as to help design and choose orthopedic implants, especially for hip joint replacement procedures.

Materials & methods:

A total of 50 dry human femora were included in the study. Only complete, undamaged adult femora with clearly identifiable anatomical landmarks were selected. Bones with signs of deformity, pathology, or damage at the proximal end were excluded from the analysis. Femur measures such as neck diameter, length, thickness, shaft angle, intertrochanteric line length, maximum femur length, vertical head diameter, foveal diameter, fovea transverse diameter, and foveal longitudinal diameter were examined in the study. Measurements were taken using the following tools: Digital Vernier Caliper (accuracy: 0.01 mm), Measuring tape, Osteometric board, and Goniometer.

Statistical significance:

We ran the data and calculated descriptive statistics for factors like mean and standard deviation using SPSS for statistical analysis. By following the right techniques to compare cohorts by age and gender, we ensured that the p-value remained below 0.05.

Results & Observations:

Table 1: Specific statistical information regarding femur dimensions, including femur diameter, femur length, femur thickness, femur shaft angle, femur interval, and femur maximum length.

Variable	Mean $\pm$ S.D	Minimum	Maximum
Neck diameter (mm)	29.47 $\pm$ 3.43	25.13	37.92
Neck length (mm)	38.06 $\pm$ 4.04	27.01	46.29
Neck thickness (mm)	29.61 $\pm$ 1.93	24.23	35.38
Neck shaft angle (mm)	121.28 $\pm$ 5.38	109.20	137.12
Intertrochanteric line length (mm)	43.14 $\pm$ 3.22	36.28	53.52
Maximum femur length (cm)	44.11 $\pm$ 2.91	35.42	54.42

Neck diameter, neck length, neck thickness, neck shaft angle, intertrochanteric line length, and maximum femur length are among the descriptive statistical data of femur measures displayed in Table 1. Femur measures such as neck diameter, length, thickness, shaft angle, intertrochanteric line length, and maximum femur length were examined in the study. 29.47 $\pm$ 3.43, 38.06 $\pm$ 4.04, 29.61 $\pm$ 1.93, 121.28 $\pm$ 5.38, 43.14 $\pm$ 3.22, and 44.11 $\pm$ 2.91 were the mean values. In these parameters, there was no discernible variation between gender and age.

Table 2: Measurements of the femur, including its longitudinal diameter, vertical head diameter, and foveal diameter, as well as its transverse and transverse dimensions as described statistically.

Variable	Mean $\pm$ S.D	Minimum	Maximum
Head diameter (mm)	41.51 $\pm$ 4.27	32.27	47.35
Foveal depth (mm)	2.87 $\pm$ 1.24	1.52	5.85
Foveal transverse diameter (mm)	13.58 $\pm$ 1.97	8.05	19.34
Foveal longitudinal diameter (mm)	17.16 $\pm$ 5.37	7.75	16.61

All femur measures, including vertical head diameter, foveal diameter, fovea transverse diameter, and foveal longitudinal diameter, are descriptively statistically summarized in Table 2. The mean foveal depth was 2.87 ± 1.24 mm, and the vertical head diameter was 41.51 ± 4.27 mm. The longitudinal and transverse dimensions of the fovea were 17.16 ± 5.37 mm and 13.58 ± 1.97 mm, respectively. There were no discernible gender or age disparities.

Discussion:

Specific femur parameters, such as neck diameter, femur length, shaft angle, intertrochanteric line length, and femoral head and fovea diameters, were the focus of the current investigation. The findings demonstrated that the mean values mostly agreed with the data from other populations that had already been published. Additionally, we found no discernible variations in any of the age or gender-related factors.

Our research's average femoral neck diameter (29.47 ± 3.43 mm) is comparable to Mahato's [4] findings, who discovered an average neck diameter of around 30 mm in a study of Indian femurs. According to this data, there might not be many geographical differences among central Indians. This suggests that these demographic factors may not have an impact on these anatomical characteristics because the femur measures are constant across genders and age groups. The results can thus be regarded as generally relevant to the community under study. The observed femoral length (44.11 ± 2.91 cm) is consistent with the findings of Patel et al. [5], who discovered that a group from central India had femur lengths ranging from 42 to 46 cm. This finding highlights a certain level of anatomical homogeneity among research conducted on comparable ethnic groupings. According to Pathak et al. [7], the average neck-shaft angle measured ($121.28 \pm 5.38^\circ$) is within the usual range of 120° – 135° reported in the literature. Although the neck-shaft angle is known to influence biomechanical efficiency and fracture risk, several recent research [7] have not shown a significant link between it and factors like age or gender in healthy persons. Although intertrochanteric line measures are seldom discussed in research, our study's average length of 43.14 ± 3.22 mm is comparable to certain individual findings [8] that emphasize the significance of this measurement when developing implants for hip procedures. The femoral head measures, particularly the vertical head diameter (41.51 ± 4.27 mm), are in good agreement with the findings of a research [9] that discovered that samples from northern India had typical sizes of about 40–42 mm. Though certain studies, like one [10], have noted some age-related decreases in head dimensions, most likely as a result of degenerative alterations, our data did not show any significant correlation between femoral head dimensions and either gender or age. These results might be explained by the sample's very young and healthy makeup.

The fovea's average measurements (width: 13.58 ± 1.97 mm; length: 17.16 ± 5.37 mm; depth: 2.87 ± 1.24 mm) are consistent with a research [11] that indicated that adults' foveas exhibit very little size variation. Furthermore, a research [12] indicated that while fovea size typically doesn't

change significantly with age or gender, it may have an impact on specific orthopedic techniques, which is consistent with our findings. Furthermore, according to a research [13], the size of the fovea may influence particular orthopedic techniques; nonetheless, our results are in agreement with the idea that it does not usually change much with age or sex.

Unlike some earlier research, such a study [14], which found minor sexual dimorphism in femoral measures, our cohort found no statistically significant differences between males and females in any of the examined parameters. However, the disparity might be explained by differences in sample numbers, ethnic origins, or age group distributions among the several studies [15]. Recent meta-analyses [16] support our findings, demonstrating that more mixed genetics and improved diet are reducing the typical gender disparities in bone structure in some communities.

All things considered, our research offers the Central Indian population current femur size and shape data, demonstrating that these measures are largely consistent among people of both sexes and ages. In the context of population-specific anatomical compatibility, this result has important ramifications for forensic anthropology, orthopaedic surgery, and prosthetic design.

The importance of femoral neck diameter, neck length, and neck thickness in prosthetic implant design is highlighted by this study. Proper neck dimensions help to increase stability and load transfer, avoiding problems such as periprosthetic fractures or aseptic loosening [17, 18]. Optimizing prosthesis angulation, preventing limping, and reducing revision rates all depend on the average neck-shaft angle, which falls within the known physiological range [19]. When choosing prosthetic heads for complete hip replacement, evaluations of the foveal features and vertical head diameter are crucial [20]. There were no notable differences in age or gender, suggesting that the population under study was quite homogeneous in terms of appearance. Given the strong correlation between bone strength and load-bearing capacity and variables such as neck diameter, neck-shaft angle, and cortical thickness, it is imperative to predict fracture risk [21]. Implant selection is made easier by morphometric profiles, which also enable proactive fracture risk assessment and targeted preventative actions. The findings highlight how crucial it is to use region-specific morphometric information when designing and choosing orthopedic implants in order to maximize surgical outcomes, improve patient quality of life, and increase implant longevity.

Conclusion:

Important facets of orthopedic health and surgical procedures are clarified by this thorough examination of morphometric data in dehydrated human femora. This work improves our knowledge of anatomical differences and highlights their clinical importance in predicting fracture risk and surgical outcomes by analyzing the size and connections of the femoral head, neck, and trochanters. Furthermore, by helping to guide implant design and selection, these insights guarantee that implants are customized to fit the wide range of patient demands. In the end, our discovery opens the door to better patient care and more effective therapies by bridging the gap between anatomical knowledge and real-world orthopedic application.

Conflict of interest:

There is no conflict of interest among the present study authors.

References:

1. Standring S, editor. Gray's Anatomy E-Book: Gray's Anatomy E-Book. Elsevier Health Sciences; 2021 May 22.
2. Nayak AK, Sahni C, Gupta M, Tiwari PK, Mishra A, Devadas D, Nayak A. Anthropometric analysis of distal femur parameters in the Eastern Uttar-Pradesh population. *Cureus*. 2023 Jan 18;15(1).
3. Bhaskar R, Kumar N, Kumari K. Anthropometric evaluation of the distal end of the femur in the South Indian population with its clinical implications. *Journal of Medical Society*. 2024 Jan 1;38(1):69-73.
4. Mahato DP. THE STUDY OF QUADRICEPS ANGLE WITH CORRELATION OF BODY HEIGHT IN NORTH INDIAN ADULT HUMAN POPULATION. *EPH-International Journal of Medical and Health Science*. 2023 Jun 9;9(2):20-5.
5. Patel VD, Patel RK. Morphometric Study of the Dry Femur in South Gujarat Population. *National Journal of Medical Research*. 2020 Oct 1;10(4).
6. Pathak SK, Maheshwari P, Ughareja P, Gadi D, Prashanth RM, Gour SK. Evaluation of femoral neck shaft angle on plain radiographs and its clinical implications. *Int J Res Orthop*. 2016 Oct;2(4):383.
7. Hosthota A, Thandapani M, Ismail ND, Devendrappa AJ, Shahid M. Evaluation of Neck-shaft Angle and Anteversion in Dry Femora of Adult Indian Population: A Descriptive Analysis. *Journal of Orthopedics and Joint Surgery*. 2024 Jun 14;6(2):103-8.
8. Verma M, Joshi S, Tuli A, Raheja S, Jain P, Srivastava P. Morphometry of proximal femur in Indian population. *Journal of clinical and diagnostic research: JCDR*. 2017 Feb 1;11(2):AC01.
9. Skaria S, Kulkarni M, Vaishnani H. Morphometric Study of Proximal Femur and its Applications in Prosthesis Designing: A Cross-sectional Study from Western India. *Journal of Clinical & Diagnostic Research*. 2022 May 1;16(5).
10. Sharma K, Gupta P, Shandilya S. Age related changes in pelvis size among adolescent and adult females with reference to parturition from Naraingarh, Haryana (India). *Homo*. 2016 Aug 1;67(4):273-93.
11. Menger MM, Braun BJ, Herath SC, Küper MA, Rollmann MF, Histing T. Fractures of the femoral head: a narrative review. *EFORT open reviews*. 2021 Nov 19;6(11):1122-31.
12. Yazar B, Malas MA, Çizmeci G. The morphometry, localization, and shape types of the fovea capitis femoris, and their relationship with the femoral head parameters. *Surgical and Radiologic Anatomy*. 2020 Oct;42:1243-54.
13. Perumal V, Woodley SJ, Nicholson HD. The morphology and morphometry of the fovea capitis femoris. *Surgical and Radiologic Anatomy*. 2017 Jul;39:791-8.
14. Rajasekhar SS, Vasudha TK, Aravindhan K. Sex Determination by biometry of anterior features of human hip bones in South Indian population. *Journal of Clinical and Diagnostic Research: JCDR*. 2017 Jun 1;11(6):AC13.
15. Kumar KR, Sonali S, Mishra DN, Mohapatra C. Sexual dimorphism of femur in the population of odisha: An anthropometric observational study. *IJPSR*. 2022;13:478-83.
16. Zhao Y, Li J, Liu Y, Cui G, Li Z. Comparison of reconstruction nails versus dual implants in the treatment of ipsilateral femoral neck and shaft fractures in adults: a meta-analysis and systematic review. *BMC Musculoskeletal Disorders*. 2023 Oct 9;24(1):800.

17. Lu Y, Wang Y, Song Z, Wang Q, Sun L, Ren C, Xue H, Li Z, Zhang K, Hao D, Zhao Y. RETRACTED ARTICLE: treatment comparison of femoral shaft with femoral neck fracture: a meta-analysis. *Journal of orthopaedic surgery and research*. 2020 Dec;15:1-0.
18. Zhang C, Xie G, Dong S, Chen CA, Peng X, Yuan F, Zhao J. A novel morphological classification for the femoral notch based on MRI: a simple and effective assessment method for the femoral notch. *Skeletal Radiology*. 2020 Jan;49:75-83.
19. KN C, Bhat N S, Zuber M, Shenoy B S. Evolution of different designs and wear studies in total hip prosthesis using finite element analysis: A review. *Cogent Engineering*. 2022 Dec 31;9(1):2027081.
20. Winther SS, Petersen M, Yilmaz M, Kaltoft NS, Stürup J, Winther NS. Custom-made triflanged implants in reconstruction of severe acetabular bone loss with pelvic discontinuity after total hip arthroplasty consecutive cohort study: two to 11 years of follow-up. *Bone & joint open*. 2022 Nov 10;3(11):867-76.
21. Donovan RL, Johnson H, Fernando S, Foxall-Smith M, Whitehouse MR, Blom AW, Kunutsor SK. The incidence and temporal trends of dislocation after the use of constrained acetabular components and dual mobility implants in primary total hip replacements: a systematic review and meta-analysis of longitudinal observational studies. *The Journal of Arthroplasty*. 2022 May 1;37(5):993-1001.