



Research Article

STUDY OF UV MUTATED PROTEASE ENZYME FROM VARIOUS SOURCES

K. Saranya *, M. Dhanachezhian, G. Gopinath, A. Mathan kumar, N.G. Ramesh babu

Department of Biotechnology, Adhiyamaan College of Engineering, Hosur, India

*Corresponding Author Email: ksaranya31@gmail.com

Article Received on: 30/11/18 Approved for publication: 11/02/19

DOI: 10.7897/2230-8407.1005167

ABSTRACT

Proteases have wide variety of functions and commercial applications. Protease is the third largest groups of industrial enzymes and is used in detergents production, leather industry, food industry, pharmaceutical industry and bioremediation processes. Protease and amylase are well known industrially important enzymes. To study the UV mutated protease enzyme from various sources. Microbes were isolated from curd, buccal cavity are probiotics were maintained under optimum conditions, the microbial growth and enzyme production carried out at 37 °C for at 24 hours and pH 7.0. The isolated microbes were mutated using UV radiation exposure which enhanced protease and amylase enzymes production. The protease enzyme was confirmed using protease and amylase assay. The isolated bacteria have the ability to produce the industrially valuable enzymes which have various industrial applications. The protease enzyme has wide applications in many industries and therefore the UV mutated enzyme will have more thermostability.

Keywords: amylase, protease, UV radiation, mutation

INTRODUCTION

Proteases have wide variety of functions and commercial applications¹. Protease is the third largest groups of industrial enzymes and is used in detergents production, leather industry, food industry, pharmaceutical industry and bioremediation processes^{2,3}. They are classified into various groups such as alkaline protease, serine protease, cysteine protease, aspartic protease and metallo protease⁴. Currently 13 billion tons of alkaline proteases find their applications in house hold laundry as detergents. Alkaline proteases were the first enzyme produced in bulk. Plant, animal and microbial sources are employed in there enzyme production. Microbial proteases are preferred to plant and animal sources to various advantages. A variety of microorganisms such as bacteria, fungi, yeast and Actinomycetes are known to produces enzymes⁵. Among all the microorganisms, *Bacillus* species produces a wide variety of extra-cellular enzymes like protease, amylase and lipase^{6,3} e.g. *B. cereus*, *B. stercorarius*, *B. mojavensis*, *B. megaterium* and *B. Subtilis*.^{7,8} Mutant *B. cereus* has the ability to sporulate in the presence of a high concentration of amino acids and glucose to produced excessive amounts of extracellular protease⁹. Amylases are extracellular enzymes that have the ability to hydrolyze the glycosidic linkages of starch and glycogen. Amylase enzyme were obtained from various sources, but amylases from fungi are the most important in biotechnology and food technology applications.¹ The present study was focused for the production of curd, buccal and probiotics isolated from Trichy Research Institute for Biotechnology at Trichy in Tamil Nadu.

MATERIALS AND METHODS

Collection of samples

The curd and probiotics sample were collected from Trichy Research Institute for Biotechnology at Tamil Nadu in India and

the buccal cavity sample was collected from buccal cavity of a volunteer. The reagents and chemicals used for this project were taken from SRL India.

Isolation and Identification

Curd

A loop full of curd was taken and it was dissolved in 5 ml of Nutrient broth and incubated at 37 °C for 24 h with pH 6.8 in control incremental condition for enrichment. Subculture was prepared using spread plate technique¹⁰.

Buccal cavity

A sample was swabbed from the buccal cavity of a human volunteer and it was dissolved in 5 ml of Nutrient broth and incubated at 37 °C for 24 h with pH 6.8 in control incremental condition for enrichment plate¹¹.

Probiotics

The pre stored probiotic sample was taken from *Bacillus cereus* and mixed with 5 ml of nutrient broth. After inoculation of the sample, the mixture was incubated at 37°C for 24 h with pH 6.8. The nutrient agar was prepared and a loop full of sample was streaked in the plate by quadrant streaking method and incubated at 37 °C for 24 h¹².

Biochemical activity analysis for microbes

Grams' staining was performed to identify grams reaction¹³. Motility test was carried out for the spontaneous movement of a cell from one location to another by consumption of energy. Morphology was used to determine the shape of the bacteria. Indole test was performed on bacterial species to determine the

ability of the organism to convert tryptophan into Indole by the deamination of tryptophan. Methyl red test was used to identify bacteria producing stable acid by mechanisms of mixed acid fermentation of glucose. The above biochemical activity tests were carried out for all selected microorganisms⁷.

Mutant colony selection

The isolated pure culture was spread over the nutrient agar medium on the plate. The plate was separated into two portions; one half of the portion was covered with aluminum foil that protects microbes from the UV exposure. The plate was placed under UV exposure for different time periods of (10, 20, 30 min) and incubated at 37 °C for 24 h, mutant colony were used for future study¹⁴.

Screening for proteolytic activity

Milk solids (w/v), 1.5% agar (w/v), 0.5% beef extract (w/v), 0.3% yeast extract (w/v) were incubated at 37°C for 24 h with pH6¹⁵.

Protease production

The culture medium used in the protease production contained 0.5% glucose (w/v), 0.75% peptone (w/v), 0.5% (w/v) MgSO₄·7H₂O, 0.5% (w/v), KH₂PO₄ (w/v), and 0.01% (w/v) FeSO₄·7H₂O (w/v) maintained at 37°C for 24 h with pH 6.8 in a shaking incubator at (140 rpm). At the end of each fermentation period, the entire fermentation broth was centrifuged at 10,000 rpm at 4°C for 15 min and the clear supernatant was used for crude enzyme preparation¹⁶.

SDS Page

The SDS page technique was carried out protocol¹⁷. The protein samples such as buccal cavity, curd and probiotic extracts were subjected to 10 % SDS PAGE with 200 KDa protein markers.

RESULTS

Isolation and identification

The bacteria isolated were identified based on their morphology, growth conditions, gram stain, motility and biochemical tests [Table 1].

The probiotic microbe isolated was characterized by the cocci shape, gram negative bacteria were positive for motility, Indole, methyl red similarly the buccal cavity 2 microbe isolated was characterized by rod shape gram negative bacteria which were positive for motility, Indole, methyl red. The Buccal cavity 1 microbe isolated was characterized by the rod shape, gram positive bacteria, presence of motility, methyl red and absence of Indole. The Curd 1 microbe isolated was characterized by rod shape, gram negative bacteria, presence of methyl red, Indole, and absence of motility. The Curd 2 microbe isolated was characterized by rod shape, gram positive bacteria, and presence of Indole and absence of motility, methyl red.

Mutant colony selection

After 24 h of incubation the mutant colony was observed in UV exposure portion of the plate. The Figure 1 showed more colonies in the UV unexposed portion of the plate. In Figure 4, UV exposed portion decreases with increasing time [Figure 1] [Figure 2] [Figure 3] [Figure 4].

The microbes isolated from buccal cavity 2 was shown maximum UV exposure with stability similarly the microbes isolated from curd was 2 shown minimum UV exposure.

Screening for proteolytic activity

The zone formation represents the lysis of skim milk agar medium. A clear zone of skim hydrolysis indicates of protease producing organism *Bacillus cereus* [Figure 5].

Table 1

S. No	Microorganism	Morphology	Gram Positive / Negative	Motility	Indole	Methyl red
1	Probiotic	Cocci	Gram negative	+	+	+
2	Buccal 1	Rod	Gram positive	+	+	+
3	Buccal 2	Rod	Gram positive	+	+	+
4	Curd 1	Rod	Gram negative	+	+	+
5	Curd 2	Rod	Gram positive	+	+	+




Figure 1: Mutant colony selection at 10 min UV exposure




Figure 2: Mutant colony selection at 20 min UV exposure




Figure 3: Mutant colony selection at 30 min UV exposure





Figure 4: Effect of mutant colony selection under UV exposure



Figure 5: Screening for proteolytic activity of *Bacillus cereus*



1-Probiotic 1, 2-Probiotic 2, 3-Buccal 1,
4-Buccal 2, 5-Buccal soup, 6-probiotic soup

Figure 6: Conformation test for protease using SDS-PAGE

SDS-Page

The presence of protease enzyme was observed only in probiotic supernatant at 45 kDa compare to curd and buccal. Furthermore, from the present study the isolated microbes in probiotic show their presence on SDS-Page [Figure 6].

DISCUSSION

The probiotic bacteria have beneficial effects due to factors includes regulation of intestinal microbial homeostasis, changes in the availability of nutrients bring and modulation of local and systemic immune response¹⁸.

The previous study reported that protease enzyme from probiotics showed a molecular weight of 37 kDa¹⁹. A study conducted in probiotics showed a molecular weight of 30 kDa²⁰.

A study conducted in probiotics showed the three predominant protein molecules from the SDS-Page of Hi Trap ion exchange²¹. The loss of non-protein compound of protease enzymes seen with weight of 34 kDa, 17 kDa, 13 kDa. The protease enzyme was observed in 45th band.

CONCLUSION

The identified *Bacillus cereus* has the ability to produce protease enzyme. Protease enzyme shows higher activity at UV mutated conditions. The confirmation of protease enzyme was identified by SDS-Page. In industry, the protease enzyme plays a vital role in the production of detergents, leather industry, food industry, pharmaceutical industry and bio remediation processes.

ACKNOWLEDGEMENT

The authors are thankful Trichy Research Institute for Biotechnology at Tamil Nadu in India for their encouragement and support in carrying out the work in Screening and UV mutated of protease enzyme.

REFERENCES

1. Janser R, Castro S De and Sato HH. Synergistic effects of agro industrial wastes on simultaneous production of protease and α -amylase under solid state fermentation using a simplex centroid mixture design; 2013. p. 813-821.
2. Wei Wang, Rita Vignani, Monica Scali, Maura Cresti M. A universal and rapid protocol for protein extraction from recalcitrant plant tissues for proteomic analysis 2006; 27(13): 2782-2786.
3. Rao MB, Tanksale AM, Ghatge MS and Deshpande VV. Molecular and Biotechnological Aspects of Microbial Proteases. Microbiology and Molecular Biology Reviews 1998; 62(3): 597-635.
4. Balachandran C, Duraipandiyan V and Ignacimuthu S. Purification and characterization of protease enzyme from actinomycetes and its cytotoxic effect on cancer cell line 2012; (A549): 392-400.
5. A Rasheedha Banu, M Kalpana Devi, GR Gnanaprabhal, B V Pradeep, M Palaniswamy. Production and characterization of pectinase enzyme from *Penicillium chrysogenum* 2010; 3(4): 377-381.
6. Puri S, Beg QK and Gupta R. Optimization of alkaline protease production from *Bacillus* sp. by Response surface methodology. Current Microbiology 2002; 44(4): 286-290.
7. Olajuyigbe F and Ajele J. Production dynamics of extracellular protease from *Bacillus* species. African Journal of Biotechnology 2005; 4(8): 776-779.
8. Karlsson A and Arvidson S. Variation in Extracellular Protease Production among Clinical Isolates of Microbiology 2002; 70(8): 4239-4246.
9. Angelo N and Holt SC. Regulation of Extracellular Protease Production in *Bacillus cereus* T: Characterization of Mutants Producing Altered Amounts of Protease 1971; 106(3): 1016-1025.
10. Sanjeeb kumar Mandal, Sunitha Vuppu S. Preliminary studies on probiotic potential of selected lactobacillus VIT SSV strains screened from curd samples of Vellore, Bihar, Haryana and Varanasi, International Journal of Pharma and Biosciences 2013; 4(2): 193-200.
11. Hamoud Khalid Aishaya, Saleh Ahmed Alogla Khalid Sukhail G Alshammari, Carlo L Isolation and Characterization of Bacterial Species from Patients with Dental Caries and Caries-free Subjects; 2016. p. 2-6.

12. Dharmaraj Ramesh, Annadurai Vinothkanna, Amit Kumar Rai, Venkata Subramanian. Isolation of potential probiotic *Bacillus* spp and assessment of their subcellular components to induce immune responses in *Labeo rohita* against *Aeromonas hydrophila*. Fish and shellfish immunology 2015; 45(2): 268–276.
13. Mahmoud W Yaish, Irin Antony, Bernard R Glick. Isolation and characterization of endophytic plant growth- promoting bacteria from date palm tree (*Phoenix dactylifera* L.) and their potential role in salinity tolerance 2015; 107(6): 1519-1532.
14. Sreedevi Basavaraju, Chandrasekhar Kathera, Pramoda Kumari Jasti PK. Purification, characterization and application of novel alkaline protease from new *Bacillus cereus* UV -15 mutant 2017; 7(4): 1-12.
15. Dalal Rupali. Original Research Article Screening and Isolation of Protease Producing Bacteria from Soil Collected from Different Areas of burhanpur Region (MP) India 2015; 4(8): 597–606.
16. Kubendran Devaraj, Salma Aathika, Karthik Periyasamy, Premkumar Manickam Periyaraman. Production of thermostable multiple enzymes from *Bacillus amyloliquefaciens* KUB29. Natural Product and Research; 2018. p. 1-4.
17. Jennifer Vandooren, Nathalie Geurts, Erik Martens, Philippe E Van Steen Ghislain. Zymography methods for visualizing hydrolytic enzymes 2013; 10(3): 211–220.
18. MS Hossain, MAA Al Bari, Mir II Wahed. Biochemical Characterization of Probiotics Available in Bangladesh, Journal of Scientific Reserch 2016; 8(1): 101-108.
19. Orhan E, Amay D and Guvenilir Y. Partial Purification and Characterization of Protease Enzyme from *Bacillus subtilis* and *Bacillus cereus*. Applied Biochemistry and Biotechnology 2005; vol 121-124: 183-194.
20. Monika Prakash, Rathindra Mohan Banik RM, Claudia Koch - Brandt. Purification and characterization of *Bacillus cereus* Protease suitable Detergent Industry. Applied Biochemistry and Biotechnology 2005; Vol 127: 143-155.
21. Yendri Junaidi , Ambar Pertiwinigrum, Yuny Erwanto and Nanung Agus Fitriyanto Semi Purification and Identifications Molecule protein Weigh of Alkaline Protease Enzyme from *Bacillus cereus* LS2B. International Journal of Bio-Science and Bio-Technology 2016; Vol. 9(3): 89-100.

Cite this article as:

K. Saranya et al. Study of UV mutated protease enzyme from various sources. Int. Res. J. Pharm. 2019; 10(5):80-84
<http://dx.doi.org/10.7897/2230-8407.1005167>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.