



Research Article

NEMATODE *MELOIDOGYNE INCOGNITA* CHITWOOD INDUCED BIOTIC STRESS ON THE PHYTOCHEMICAL COMPOSITION OF MULBERRY LEAVES

P. Victoria Rani * and N. Vijaya Kumari

Research Scholar, Department of Sericulture, Sri Padmavathi Mahila University, Tirupati, Andhra Pradesh, India

*Corresponding Author Email: pvrani25@gmail.com

Article Received on: 28/12/17 Approved for publication: 20/01/18

DOI: 10.7897/2230-8407.09116

ABSTRACT

Root-knot disease caused by nematode *Meloidogyne incognita* Chitwood cause considerable damage to the growth and development of plants and also reduced nutritive value of leaves especially in mulberry, which is the main food source of silkworm *Bombyx mori* L. The nutritional quality of the mulberry leaf has considerable effect on the silk worm cocoon crop yield. Hence, the present investigation was carried out to know the effect of root knot nematode on important phytochemicals of mulberry leaves infested with root knot nematode *Meloidogyne incognita*. After a careful study it was found that *M. incognita* induced biotic stress has effected considerable changes in biomolecule composition as well as vital enzymatic activity in mulberry leaf and also on the total qualitative yield of the leaf.

Key words: Biomolecules, Enzymes, *Meloidogyne incognita*, Mulberry, Root knot disease.

INTRODUCTION

Mulberry (*Morus* sps) belongs to the family *Moraceae*, which is a highly valuable plant for silk industry. Its foliage forms as the main food source for growing silk worm *Bombyx mori* L. Mulberry is usually subject to a number of diseases caused by bacteria, virus, fungi, mycoplasma and nematodes. These may cause 10-30% loss in the total yield of the leaf and may reduce the nutritive values of the mulberry leaves^{1,2}. Of these, the species of nematode *Meloidogyne incognita* (Kofaid and White) Chit Wood cause a great economic loss in the yield of mulberry³. This is often found in sandy loamy soils under irrigated conditions⁴. Bessey was the first person who identifies the effect of the root-knot disease on mulberry from U.S.A⁵. The root knot nematode *Meloidogyne incognita* cause galls or knots on roots of its host plants. Formation of galls/knots on the roots and other symptoms found above ground including stunted growth, chlorosis, wilting, leaf curling and reduced vigor of the plant are the conspicuous symptoms of the disease on the mulberry plant⁶.

The perennial nature of mulberry easily succumbs to the root knot nematode once the nematode enters the field. Hence, the biotic stress of root knot nematode on the phytochemical components of mulberry plants and consequent loss of the yield have interested the researcher to take up this present study. The finding of the study are hoped to benefit the mulberry plant growers get rid of a common disease among plants and would help them avoid loss of yield.

MATERIALS AND METHODS

The study was carried out in the department of Sericulture, Sri Padmavathi Mahila University, Tirupati, Andhra Pradesh, during the period 2014-2017. For this experiment, seventy days old mulberry V1 variety saplings were planted in randomized block design with 3' x 3' spacing. After three months of establishment,

1000 juveniles/plant was inoculated as ECL (Economic Threshold Level and Crop Loss)⁷ keeping control. Sixty days after inoculation biomolecule compounds like Total chlorophylls, Carbohydrate, Protein, Starch and stress induced biomolecules (stress markers) such as Phenols, Free amino Acids, Proline, Ascorbic acid, Chlorogenic acid, Tryptophan were analyzed. In addition to that different non stress enzyme specific activity such as Malate dehydrogenase, Nitrate reductase, Amylase, Indole acetic acid oxidase (leaf and root) stress induced enzymes like Catalase, Super oxide dismutase, Peroxidase, Ascorbate peroxidase, Glutathione reductase, Polyphenol oxidase, Phenylalanine ammonia lyase, was studied in the leaves of mulberry plants.

Biomolecule estimations

Chlorophylls

The chlorophyll content in leaves was estimated by the method⁸ and expressed as mg chl/g tissue was calculated using the following formula.

$$\text{Chl} - a \text{ (mg/g tissue)} = 12.7(A_{663}) - 2.69(A_{645}) \times V / 1000 \times W$$

$$\text{Chl} - b \text{ (mg/g tissue)} = 22.9 (A_{645}) - 4.68 (A_{663}) \times V / 1000 \times W$$

$$\text{Total chl (mg/g tissue)} = 20.2 (A_{645}) - 8.02 (A_{663}) \times V / 1000 \times W$$

Where A = absorbance at specific wavelengths

V = final volume of chlorophyll extract in 80% acetone

W = fresh weight of tissue extracted

Carbohydrate

The total carbohydrate was estimated by anthrone method⁹. Carbohydrate was calculated in relation to fresh weight basis and expressed as mg/g. The calculation is done by using the formula.

Amount of carbohydrate present in 100 mg of the sample

$$= \frac{\text{Mg of glucose}}{\text{Volume of test sample}} \times 100$$

Protein

Protein content of leaves was estimated by Lowry's method¹⁰. The amount of protein mg/g sample was expressed.

Starch

The starch content was estimated by following the method⁹ and the starch content was expressed as mg/g of leaf.

Stress induced biomolecules

Phenols

Phenols were estimated following the method¹¹. Concentration of phenols was expressed in $\mu\text{g/g}$ fresh weight material, equivalent to catechol.

Free amino acids

Free amino acids were estimated following the method¹². Computation was done using standards and the amount of amino acids was expressed as $\mu\text{g/g}$.

Proline

Proline estimation was done by the following method¹³ and expressed in $\mu\text{g/g}$ fresh weight.

$$\mu\text{g/g} = \frac{\mu\text{g proline /ml} \times \text{ml toluene}}{115.5 \times \text{g of sample}} \times 5$$

Where 115.5 is the molecular weight of proline.

Ascorbic Acid

Estimation of Ascorbic acid was followed by the method¹⁴ and expressed as mg/g.

Chlorogenic acid

Estimation of chlorogenic acid was followed by the method¹⁵. Chlorogenic acid content was calculated of the sample from the standard graph and it was mentioned as mg/g.

Tryptophan

The Tryptophan content was estimated by following the method¹⁶. It was calculated by using the following formula and which was expressed as $\mu\text{g/g}$.

$$\text{Tryptophan in leaf sample} = \frac{\text{Tryptophan value from graph in } \mu\text{g}}{\text{Percent N in the sample}} \times 0.096$$

Leaf Yield / ha/ annum (Kg)

Leaf yield was calculated on the basis of spacing given for plantation, leaf yield/ hectare/ annum was calculated by using the formula.

$$= \frac{\text{Total number of plants in an acre} \times 2.5 \times \text{Total yield}}{\text{plant} \times \text{Total number of crops / annum.}}$$

Enzyme activity in Mulberry leaves

Non stress enzymes

Nitrate reductase

The enzyme activity was studied following the method¹⁷. Enzyme activity is expressed as micromole nitrite produced $\mu\text{M min}^{-1}\text{mg}^{-1}$ protein.

Malate dehydrogenase

The enzyme activity was assayed by the following method¹⁸. The enzyme activity was expressed as $\text{min}^{-1}\text{mg}^{-1}$ protein.

Amylase Activity

The amylase inhibition was studied by the method¹⁹ and expressed as $\text{unit min}^{-1}\text{mg}^{-1}$ protein.

Stress induced enzymes

Catalase activity

Catalase activity was estimated following the method²⁰. Enzyme activity was computed by calculating the amount of H_2O_2 decomposed and expressed as $\text{unit min}^{-1}\text{mg}^{-1}$ protein.

$$\text{Unit activity} = \frac{\text{Change in Abs / min} \times \text{total volume (ml)}}{\text{Ext. coefficient} \times \text{volume of sample used (ml)}}$$

Where Ext. coefficient = $6.93 \times 10^{-3}\text{mM}^{-1}\text{cm}^{-1}$

Superoxide dismutase activity

SOD was assayed following the method²¹. Unit of enzyme activity was expressed as enzyme unit (EU) $\text{min}^{-1}\text{mg}^{-1}$ protein.

$$\text{Unit of enzyme} = \frac{\text{Blank} - \text{Sample}}{\text{Blank}/2}$$

Ascorbate Peroxidase

Ascorbic acid was assayed by following the method²². Enzyme activity was calculated as concentration of ascorbic acid oxidized and expressed in $\text{unit min}^{-1}\text{mg}^{-1}$ protein.

$$\text{Unit activity} = \frac{\text{Change in Abs / min} \times \text{total volume (ml)}}{\text{Ext. coefficient} \times \text{volume of sample used (ml)}}$$

Where Ext. coefficient = $2.8 \times 10^{-3}\text{mM}^{-1}\text{cm}^{-1}$

Peroxidase

Peroxidase activity was determined specifically with Guaiacol at 436 nm following the method²³. The enzyme activity was expressed as $\text{unit min}^{-1}\text{mg}^{-1}$ protein.

$$\text{Enzyme activity units/litre} = \frac{3.18 \times 0.1 \times 1000}{6.39 \times 1 \times \Delta t \times 0.1}$$

Glutathione reductase (Gr)

Glutathione reductase activity was determined by following the method²⁴. The enzyme activity was expressed as $\text{min}^{-1}\text{mg}^{-1}$ protein.

Phenylalanine ammonia lyase (Pal)

Enzymatic assay of Phenylalanine ammonia lyase (PAL) activity was analyzed as the rate of conversion of L-phenylalanine into

trans-cinnamic acid at 270 nm in a spectrophotometer as mentioned²⁵ and was expressed in $\text{min}^{-1} \text{mg}^{-1}$ protein.

$$\text{Units/ml enzyme} = \frac{(\Delta A_{270\text{nm}}/\text{min Test} - \Delta A_{270\text{nm}}/\text{min Blank}) (3) (\text{df})}{(19.73) (0.1)}$$

Poly phenol oxidase (Ppo)

Polyphenol oxidase was assayed following method²⁶ and expressed as unit $\text{min}^{-1} \text{mg}^{-1}$ protein.

$$\text{Units in the test} = K \times (\Delta A/\text{min})$$

Where, K is 0.272 for Catechol oxidase

Indole acetic acid oxidase

IAAO activity was determined by the method²⁷. The enzyme activity was expressed as m mole IAA oxidized per $\text{min}^{-1} \text{mg}^{-1}$ protein.

Statistical analysis

All results were subjected to statistical analysis. Data was analyzed by analysis variance T-test and tested for significance at 0.01 and 0.05 levels following the method²⁸.

RESULTS AND DISCUSSION

Effect of root knot nematode *M. incognita* on biomolecule composition and stress induced biomolecules in mulberry leaves

Plant parasitic nematode, *Meloidogyne incognita* caused considerable changes in biomolecule composition of the infested mulberry leaves. Results were presented in the Table 1 and 2.

Biomolecule compounds

A considerable changes in total chlorophyll content as well as chlorophyll-a and b, carbohydrate, protein, starch and leaf yield were observed in nematode infested leaves compared to healthy plants. The significant reductions were observed in all the above parameters against healthy plants. The results are presented in Table 1 and Figures 1, 3.

Plant parasitic nematode, *Meloidogyne incognita* alters the metabolic processes of the host plant which are manifested in the form of physiological and biochemical changes occurring in the infested plants.

Hyperplasia and hypertrophy caused due to root knot disease on mulberry roots results in reduced uptake and transport of water and nutrients. Reduced water and nutrients absorptions make the plant suffer from decreased biomolecule synthesis ultimately reduces the yield. Root-knot nematodes cause giant cells to form in the roots, and this disrupts the root vascular system, reducing the uptake of water and nutrients and their transport from the roots to the shoots²⁹.

The present observations are in conformity with earlier reports. This was reported that protein and carbohydrates as well as chloroplast pigments are adversely affected due to root knot nematode *M. javanica* infestation in chickpea (*Cicer arietinum* L.)³⁰. The starch level decreased in the host plants infested by *Meloidogyne spp* as observed³¹.

Stress induced biomolecules

When nematode infested mulberry leaves were analyzed for the presence of stress induced biomolecules, there was significant increase in the percentage of Phenols, Free amino acids, Proline, Ascorbic acid, Chlorogenic acid and Tryptophan.

High concentrations of Phenols, Free amino acids, Ascorbic acid, Chlorogenic acid and Tryptophan involve in the plant defense mechanism and create unfavorable conditions for the growth of pathogen and thus induce disease resistance in host plant. This might be the reason for increased amount of stress induced biomolecules.

The observations are in conformity with earlier reports that revealed that the *M. incognita* infested okra plants showed increased levels of phenolic compounds over the healthy plants³². It was investigated the biochemical alternation resulting from *Meloidogyne incognita* infestation in *Cicer arietinum* and reported an increased level of proline, free amino acids and total phenols in infested plants³³. This was proposed that enzymes of root-knot nematodes liberate tryptophan from protein complexes and this compound converted to Indole acetic acid (IAA) could be responsible for galling³⁴.

Enzymes activity in leaves

Effect of root knot nematode, *Meloidogyne incognita* in enzyme activity of mulberry leaves.

Root knot nematode *Meloidogyne incognita* cause considerable changes in various enzyme activities in the nematode infested mulberry plants.

Non stress enzymes

Nematode infestation in mulberry plant cause reduction in Nitrate reductase and increased Malate dehydrogenase, Amylase activity and Indole acetic acid oxidase which are the major synthetic and catabolic enzymes. The reason could be the damaged root system results reduced uptake of water and nutrients. The results are presented in Table -2 and Figure-2.

Stress induced enzymes

When plants expose to any type of stress they react and try to protect them from oxidative stress which is the primary reaction. Enzymatic antioxidants like Catalase, Super oxide dismutase, Peroxidase, Ascorbate peroxidase, Glutathione reductase, Polyphenol oxidase, Phenylalanine ammonia lyase, are commonly present in the subcellular compartments which show increased activity to protect the plant system from oxidative stress. In the present study also significance increase was observed in all the stress enzyme activities. The results are presented in Table -2 and Figure-2.

Nematode induced biotic stress which leads to overproduction of Reactive oxygen species (ROS) continuously as byproducts of various metabolic pathways. They cause damage to proteins, lipids, carbohydrates and DNA resulting in oxidative stress³⁵. In favor to limit oxidative damage under stress, plants have enhanced a series of detoxification systems that break down the highly toxic ROS³⁶.

Enzyme activity is one of the important tools to confirm the resistance to root pathogenic nematodes. When a nematode infests the host tissue, a small number of specific genes are induced to produce mRNAs that permit synthesis of similar number of specific proteins³⁷. Many of these proteins are

enzymes such as phenylalanine ammonia lyase, polyphenol oxidase, peroxidase and β -1-3-glucanase. These are involved in the synthesis of low molecular weight substances such as phytoalexins, phenols and lignin which are inhibitory to the invading nematodes³⁸.

The results of the present investigations are in agreement with the various reports. The increased percentage of peroxidase activity, phenol content, polyphenol oxidase activity, phenylalanine

ammonia lyase was found in roots of egg plant infested with nematode³⁹. The investigations carried out to establish a link between the activities of catalase, peroxidase and superoxide dismutase in tomato and nematode *M. incognita* parasite interactions revealed that, increase in the enzyme activity was recorded highest after nematode inoculation for peroxidase, superoxide dismutase and catalase respectively in the susceptible cultivar⁴⁰.



Mulberry garden



Root knot diseased plant

Table 1. Effect of root knot nematode *M. incognita* on Biomolecules of mulberry leaves

S. No	Parameters	Control	Infested	% of decreased/ increased	T- Test	Significant at 0.01 level
1	Total chlorophylls (mg/g)	2.797	1.890	32.427	51.403	**
	Chl- a (mg/g)	1.860	1.200	35.483	34.467	**
	Chl- b (mg/g)	0.940	0.690	26.595	21.651	**
2	Carbohydrate (mg/g)	49.000	31.330	36.061	496.484	**
3	Protein (mg/g)	59.000	40.660	31.084	589.876	**
4	Starch (mg/g)	17.400	10.500	39.655	225.856	**
5	Leaf yield / ha/ annum (kg)	16036.370	11616.000	27.564	193.038	**
6	Phenols (μ g/g)	2.660	4.100	54.135	23.568	**
7	Free amino acids (μ g/g)	13.000	23.330	79.461	565.797	**
8	Proline(μ g/g)	3.570	5.250	47.058	49.903	**
9	Ascorbic acid (mg/g)	82.997	92.660	11.642	346.497	**
10	Chlorogenic acid (mg/g)	42.660	56.660	32.817	588.188	**
11	Tryptophan (μ g/g)	0.457	0.787	72.21	17.501	**

Note – ** Significant at 0.01 level

Table 2. Effect of root knot nematode, *M. incognita* on enzymatic activity of mulberry leaves

S.No	Parameters ($\text{min}^{-1} \text{mg}^{-1} \text{protein}$)	Control	Infested	% of decreased/ increased	T- Test	Significant at 0.01 level
1	Nitrate reductase	5.87	4.36	25.724	40.356	**
2	Malate dehydrogenase	0.337	0.563	67.062	11.031	**
3	Amylase	1.65	2.37	43.636	37.601	**
4a	Indole acetic acid oxidase in leaf	0.113	0.187	65.486	7.778	**
4b	Indole acetic acid oxidase in root	0.803	1.59	98.007	11.600	**
5	Catalase	7.723	9.057	17.273	33.806	**
6	Peroxidase	0.34	0.61	79.411	33.068	**
7	Ascorbate Peroxidase	1.163	2.01	72.828	35.223	**
8	Glutathione reductase	0.083	0.133	60.24	4.009	*
9	Superoxide dismutase	0.027	0.047	74.074	4.243	*
10	Polyphenol oxidase	0.34	0.66	94.117	24.787	**
11	Phenylalanine ammonia lyase	0.807	1.443	78.81	33.764	**

Note – ** = Significant at 0.01 level

*= Significant at 0.05 level




Figure 1. Effect of root knot nematode *M. incognita* on biomolecules of mulberry leaves




Figure 2. Effect of root knot nematode, *M. incognita* on enzyme activity of mulberry leaves




Figure 3. Effect of root knot nematode, *M. incognita* on leaf yield of mulberry

CONCLUSION

It is concluded that root-knot nematode *Meloidogyne incognita* bring about considerable phytochemical changes in infested mulberry plants may be considered as evidence of altered physiology and growth of plants due to host-parasite relationship and interaction resulting to loss in crop yield and also the increased stress enzyme activity could be resistant of plant defense mechanism against root-knot nematode *Meloidogyne incognita*.

ACKNOWLEDGMENT

The author is thank full to U.G.C for sanctioning F.I.P (Faculty Improvement Program) to do this research work and also great full to supervisor and staff members of Sericulture department, Sri Padmavati Mahila University, Tirupati. for providing facilities to carryout research work.

REFERENCES

- Chanturiya ,N.N., Labakhua, L.V. (1963) Listing of antibiotics to control bacterial infection in mulberry trees. Soobsch Akad. Nauk Gruz SSR, 32(1), 141- 148.
- Shree, M.P.Umeshkumar, N.N. (1991). Changes in the elemental composition of mulberry leaves (*Morus alba* L. var S-54) infected by foliar pathogens, Sericologia, 31(3), 441 - 444.
- Chitwood B.G. (1949). Root knot nematodes, part1. A revision of the genus *Meloidogyne* Goelidi, 1887. Proc. Helminth Soc. Wash., 16: 90-104.
- Narayanan, E.S, Kasiviswanathan, K. and Iyengar, M.N.S. 1966. Effect of varietal feeding, irrigation levels and nitrogen fertilization on larval development and cocoon characters of *Bombyx mori* L. Ibid. 5:13-17.
- Bessey, E. A.1911.Root-knot and its control. United States Dept. Agric., Bur. Plant Ind. Bull. 217. 89 pp.
- Saha , S.S., Sinhababu , S.P. and Sukul , M.C. (1983).The effect of nematodes infestation on mulberry plants and their effect on feeding silkworm *Bombyx mori*. L. Nematologica 29(4):463 -467.
- Govindaiah, Dandin, S.B. and Sharma, D.D. (1991) pathogenicity and avoidable leaf yield loss due to *Meloidogyne incognita* in mulberry (*Morus alba* L.). Indian Journal Nematol., 21: 52 -57.
- Arnon, D.I. 1949. Copper enzymes in isolated chloroplasts, polyphenol oxidase in *Beta vulgaris*. Plant Physiology, 24: 1-15.
- Hodge, J.E. and Hofreiter, B.T. (1962). In: Methods in Carbohydrate Chemistry, Vol.17(Eds.,) Whistler, R.L. and Be Miller, J.N., Academic Press, New York. p. 420.
- Lowry OH, Rosenbrough, NJ, Farr AL, Randal RJ, protein measurement with the Folin Phenol Reagent. Journal of Biological Chemistry, 193:265- 275,(1951).
- Malik E.P., Singh M.B.,(1980). Plant Enzymology and Histochemistry (1st Edn.) Kalyani Publishers, New Delhi, pp:286
- Theymoli Balasubramanian and Sadasivam. S (1987) Plant Food, Hum. Nutr. 37 P. 41.
- Bates LS, Waldren RP, Teare ID (1973) Rapid determination of free proline for water stress studies. *Plant Soil* 39: 205-207.
- Sadasivam, S and Theymoli Balasubramanian (1987) In : Practical manual in biochemistry Tamil Nadu, Agricultural university, Coimbatore p 14.
- Michael Eskin, M A, Hochn, E and Frenkel, C (1978) Journal of Agricultural and Food Chemistry, 26: 973.
- Mertz, E T, Jambunathan, R and Misra, P S (1975) In: Protein Quality Agricultural Research Station Bull No. 7, Purdue Univ., USA, p. 9.
- Sadasivam, S. and G. Gowri. 1981. Ammonia assimilatory enzymes in a nif⁺ mutant of *Azotobacter chroococcum*. Experientia, 37: 552-553.
- Sadasivam S and Gowri G (1981) Activities of phosphoenolpyruvate carboxylase, NAD⁺-malate dehydrogenase and aspartate amino-transferase and chlorophyll a/b ratio in leaves of sorghum cultivars during plant growth, Photosynthetica. 15, 453-456.
- Bernfeld, Amylase, α and β , in Methods in Enzymology, S. P. Colowick and N. O. Kaplan, Eds., pp. 149-158, Academic Press, New York, NY, USA, 1955.
- Aebi H. In: Catalase in Vitro. Methods in Enzymology. Colowick SP, Kaplan NO, editors. Vol. 105. Florida Academic Press, New York 1984. pp. 114-121.
- Dhindsa, R.H., R. Plumb-Dhindsa and T.A. Thorpe (1981). Leaf senescence correlated with increased level of membrane permeability, lipid peroxide tior and decreased level of SOD and CAT. Journal of experimental Botany, 32, 93-101.
- Nakano, Y and Asada, K. (1981). Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. Plant Cell Physiol., 22:867 -880.
- Putter, J (1974) In : *Methods of Enzymatic Analysis 2* (Ed Bergmeyer) Academic Press New York p 685.
- Smith, I.K., Vierheller, T. L., Thorne, C. A. (1988). Assay of glutathione reductase in crude tissue homogenates using 5, 5'-dithiobis (2- nitrobenzoic acid). *Anal. Biochem.*, 175: 408 -413.
- Dickerson DP, Pascholati SF, Haagerman AE, Butler LG, Nicholson RL. Phenylalanine ammonia-lyase and hydroxy l cinnamate: CoA ligase in maize mesocotyls inoculated with *Helminthosporium maydis* or *Helminthosporium carbonum*. Physiology Plant Pathology. 1984; 25: 111-123.
- Esterbaner , H. Schwarzl, E and Hayn, M (1977). A rapid assay for catechol oxidase and laccase using 2-nitro-5-thiobenzoic acid. *Anal Biochem* 77: 486- 494.
- Byrant, S.D and Lane, FE (1979) Indole-3- acetic acid oxidase from peas. *Plant Physiol* 63: 696- 699 .
- Rangaswamy, R (2000). *A text book of agricultural statistics*, New Delhi: New Age International (P) Limited Publishers.
- Abad, P., Favery, B., Rosso, M. and Castagnone-Sereno, P. 2003. Root-knot nematode parasitism and host response: molecular basis of a sophisticated interaction. Molecular Plant Pathology, 4: 217-224.
- Upadhyay KD and Banerjee B. 1986. Some chemical changes in chickpea plants infected with root-knot nematode, *Meloidogyne javanica*, Indian Journal Nematology, 16 (2): 286-288.
- Orion, D and R. Bronner. 1973. The localization of starch, amylase and invertase in *Meloidogyne javanica* galls. Nematologica., 19:401-402.
- Agarwal M, Goel, AK, Satish k and Tayal MS. 1985. Biochemical changes in okra infected with root knot nematode, *Meloidogyne incognita*, same hydrolysing and oxidative enzymes and related chemical metabolites, Indian Journal of Nematology, 15(2): 255-256.
- Sharma NT and Trivedi PC. 1987. Biochemical alterations resulting from *M. incognita* infection in *Cicer arietinum*, Indian Journal of Mycology and Plant Pathology, 17 (2): 160-164.
- Sayre, R.M. 1958. Plant tissue culture as a tool in the study of the physiology of the root-knot nematode, *Meloidogyne incognita*. Ph.D. diss., Univ. of Nebraska, Lincoln. 40 p.
- Sreedhar M, Chaturvedi A, Aparna M, Kumar PD, Singhal RK & Venu BP, Influence of γ - radiation stress on scavenging enzyme activity and cell ultra structure in groundnut (*Arachis*

- hypogaea L.). Advances in *Applied Science Research*, 2013, 4(2):35-44.
36. Larkindale J & Huang B, Thermotolerance and antioxidant systems in *Agrotis stolonifera* : involvement of salicylic acid, abscisic acid, calcium, hydrogen peroxide, and ethylene. *Journal of Plant Physiol*, 161 (2004) 405.
 37. Vera, P. and Conejero, V. 1989. The induction and accumulation of pathogenesis-related P69 proteinase in tomato during citrus exocortis viroid infection and after chemical treatments. *Physiology Molecular and Plant Pathology*, 34: 323-334.
 38. Vidhyasekaran, P. 1993. Defense genes for crop disease management, genetic engineering, tissue culture. *In: Molecular Biology for Crop Pest and Disease Management*. P. Vidhyasekaran. (ed.). Daya Publishing House, New Delhi. Pp. 17-30.
 39. Xu, Xiao Ming, Xu Kun, Yu Qin and Zhang-Xiaoyan. 2008. The relationship between resistance of *Meloidogyne incognita* and phenylpropanes metabolism in roots of eggplant root-stock. *Acta Phytophylacica Sinica*, 35 (1): 43-46.
 40. Rajasekhar S.P., Ganguly A.K., Swain S.C. 1997. Quantitative changes in superoxide dismutase catalase and peroxidase with reference to resistance in tomato to *Meloidogyne incognita*. *Indian J. Nematol.* 27 (1): 79-85.

Cite this article as:

P. Victoria Rani and N. Vijaya Kumari. Nematode *Meloidogyne incognita* chitwood induced biotic stress on the phytochemical composition of mulberry leaves. *Int. Res. J. Pharm.* 2018;9(1):107-113 <http://dx.doi.org/10.7897/2230-8407.09116>

Source of support: U.G.C., India, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.